

SUPERCONDUCTIVITY NEAR THE NORMAL STATE IN A HALF-PLANE UNDER THE ACTION OF A PERPENDICULAR ELECTRIC CURRENT AND AN INDUCED MAGNETIC FIELD

YANIV ALMOG, BERNARD HELFFER AND XING-BIN PAN

ABSTRACT. We consider the linearization of the time-dependent Ginzburg-Landau system near the normal state. We assume that a constant magnetic field and an electric current are applied through the sample, which captures half of the plane, inducing thereby, a linearly varying magnetic field. In the limit of small normal conductivity we prove that if the electric current is lower than some critical value, the normal state loses its stability. For currents stronger than this critical value, the normal state is stable. To obtain this stability result we analyze both the spectrum and the pseudo-spectrum of the linearized operator. The critical current tends, in this small conductivity limit, to another critical current which had been obtained for a reduced model which neglects magnetic field effects.

CONTENTS

1. Introduction	1
2. Analytic Dilation	7
3. The Scheme of the Proof of the Main Results	9
4. Constructions of Quasimodes	12
5. Resolvent Estimates	14
5.1. Preliminary estimates	15
5.2. Projection on $(\text{span}\{\psi_0\})^\perp$	22
6. Existence, Upper and Lower Bounds	27
7. Decay of the Semi-Group	28
Appendix A. Some Results Related to Pseudospectra	31
References	32

1. INTRODUCTION

Consider a superconductor placed at a temperature lower than the critical one. If an electric current is applied through the sample it will induce a magnetic field, and as is well-understood from numerous experimental observations [22], a sufficiently strong current will force the superconductor to arrive at the normal state. If the current is then lowered, the normal state would lose stability and the sample would become superconducting again. In addition to experimental observations a similar pattern of behaviour has also been obtained theoretically by analyzing the stability of the normal state for the time dependent Ginzburg-Landau system, but with the induced magnetic field neglected [14, 3].

1991 *Mathematics Subject Classification.* 82D55, 35B25, 35B40, 35Q55.

Key words and phrases. superconductivity, critical current, critical magnetic field.

In a recent contribution [4] we analyzed the stability of the normal state in the presence of an electric current which induces a magnetic field, but in the absence of a boundary. We offered an analysis of a two-dimensional setting, i.e, in \mathbb{R}^2 , which is the simplest case where one can consider the effect of a magnetic field induced by a current, with boundary effects ignored. We found in [4] that the normal state is always stable, irrespectively of current intensity. This result is in line with those obtained for a reduced model where the magnetic field has been neglected [14, 3].

In the present contribution we introduce the effect of the boundary of the sample. With magnetic field neglected this effect has been analyzed by considering a one-dimensional problem on the half-line $\mathbb{R}_+ = \{x \in \mathbb{R} : x > 0\}$ with a Dirichlet boundary condition at $x = 0$, which stands for a normal/superconducting interface [14, 3]. Due to this effect the normal state loses its stability when the current is weaker than a certain critical value. It has been proved in [3] that the critical current for a large bounded three-dimensional domain is bounded from above by the one-dimensional value. Furthermore, for a current below the critical one, a short-time instability was proven [3] (the question whether the normal state is unstable for such domains and currents is still open).

We expect the boundary effect on the stability of the normal state in the presence of an induced magnetic field, to be similar to their effect in the absence of magnetic fields [14, 3]. To examine the boundary's impact, we consider here a problem in the half-plane \mathbb{R}_+^2 . If our expectation is correct, then in a similar fashion to [14, 3] the normal state would lose its stability for currents lower than some critical value, which we obtain here in the limit of small normal conductivity.

Assuming that a magnetic field described by \mathcal{H}^e is perpendicularly applied to the sample, the time-dependent Ginzburg-Landau system can be written as follows (see for instance [5, 6, 9, 10, 14, 21, 23]):

$$\begin{cases} \partial_t \psi + i\kappa \Phi \psi = \nabla_{\kappa \mathbf{A}}^2 \psi + \kappa^2(1 - |\psi|^2)\psi & \text{in } (0, T) \times \mathbb{R}_+^2, \\ \kappa^2 \text{curl}^2 \mathbf{A} + \sigma(\partial_t \mathbf{A} + \nabla \Phi) = \kappa \text{Im}(\bar{\psi} \nabla_{\kappa \mathbf{A}} \psi) + \kappa^2 \text{curl} \mathcal{H}^e & \text{in } (0, T) \times \mathbb{R}_+^2, \\ \psi = 0, \quad -\frac{\sigma}{\kappa^2} \frac{\partial \Phi}{\partial \nu} = J & \text{on } \partial \mathbb{R}_+^2, \end{cases} \quad (1.1)$$

where ψ is the order parameter, \mathbf{A} is the magnetic potential, Φ is the electric potential, the Ginzburg-Landau parameter of the superconductor is denoted by κ , the normal conductivity of the sample is denoted by σ , the magnitude of the dimensionless electric current is denoted by J , and the applied magnetic field is denoted by \mathcal{H}^e . In (1.1) we use the notation

$$\nabla_{\kappa \mathbf{A}} \psi = \nabla \psi - i\kappa \mathbf{A} \psi, \quad \nabla_{\kappa \mathbf{A}}^2 \psi = \Delta \psi - i[2\mathbf{A} \cdot \nabla \psi + \psi \text{div} \mathbf{A}] - |\mathbf{A}|^2 \psi,$$

where $i = \sqrt{-1}$. The half-plane \mathbb{R}_+^2 is defined in the following manner

$$\mathbb{R}_+^2 = \{(x, y) \in \mathbb{R}^2 : y > 0\}.$$

The triplet (ψ, \mathbf{A}, Φ) should also satisfy an initial condition at $t = 0$.

A solution (ψ, \mathbf{A}, Φ) to (1.1) is called a normal state solution if $\psi \equiv 0$. From (1.1) we see that if $(0, \mathbf{A}, \Phi)$ is a time-independent normal state solution then (\mathbf{A}, Φ) satisfies the following equation

$$\kappa^2 \text{curl}^2 \mathbf{A} + \sigma \nabla \Phi = \kappa^2 \text{curl} \mathcal{H}^e \quad \text{in } \mathbb{R}_+^2. \quad (1.2)$$

By taking the divergence of (1.2) we obtain

$$\begin{cases} \Delta \Phi = 0 & \text{in } \mathbb{R}_+^2, \\ \frac{\partial \Phi}{\partial \nu} = -\frac{\kappa^2 J}{\sigma} & \text{on } \partial \mathbb{R}_+^2. \end{cases} \quad (1.3)$$

Since we expect solutions of (1.3) to represent the electric potential at the normal state near the boundary of a large bounded domain, we look for solutions with bounded gradient (or $\nabla\Phi \in L^\infty(\mathbb{R}_+^2)$). Assuming that the current is of constant magnitude J along $\partial\mathbb{R}_+^2$, and that its direction is always perpendicular to it, we obtain that the unique solution to (1.3) which obeys these assumptions is given, up to an additive constant, by

$$\Phi = \frac{\kappa^2 J}{\sigma} y. \quad (1.4)$$

Assuming further that the applied magnetic field is, like the current, of constant magnitude as well, we obtain

$$\mathcal{H}^e = h \mathbf{i}_z,$$

throughout the entire sample. Here $\mathbf{i}_x, \mathbf{i}_y$ and \mathbf{i}_z denote the canonical basis in \mathbb{R}^3 . Hence, we consider an applied magnetic field which is perpendicular to the sample and parallel to its surface. Under these additional assumptions, equation (1.2) admits the following solution

$$\mathbf{A} = \frac{1}{2J} (Jx + h)^2 \mathbf{i}_y. \quad (1.5)$$

For the above \mathbf{A} and Φ , $(0, \mathbf{A}, \Phi)$ is a normal state solution of (1.1). Note that the magnetic field

$$\mathbf{H} = \text{curl}\mathbf{A} = (Jx + h) \mathbf{i}_z,$$

is the sum of the constant applied magnetic field $h \mathbf{i}_z$ and a linear term $Jx \mathbf{i}_z$ induced by the electric current.

The linearization of (1.1) near the normal state solution $(0, \mathbf{A}, \Phi)$ obtained above yields a linear equation

$$\begin{cases} \partial_t \psi + \frac{i\kappa^3 J y}{\sigma} \psi = \Delta \psi - \frac{i\kappa}{J} (Jx + h)^2 \partial_y \psi - \left(\frac{\kappa}{2J}\right)^2 (Jx + h)^4 \psi + \kappa^2 \psi & \text{in } (0, T) \times \mathbb{R}_+^2, \\ \psi = 0 & \text{on } \partial\mathbb{R}_+^2. \end{cases} \quad (1.6)$$

Applying the transformation

$$(t, x, y) \rightarrow \left(t, x - \frac{h}{J}, y\right)$$

we obtain

$$\begin{cases} \partial_t \psi + \frac{i\kappa^3 J y}{\sigma} \psi = \Delta \psi - i\kappa J x^2 \partial_y \psi - \left(\left(\frac{\kappa J}{2}\right)^2 x^4 - \kappa^2\right) \psi & \text{in } (0, T) \times \mathbb{R}_+^2, \\ \psi = 0 & \text{on } \partial\mathbb{R}_+^2. \end{cases} \quad (1.7)$$

In the present contribution we analyze the asymptotic behavior of the solutions of (1.7) for large t . We assume $J > 0$ in the sequel. Otherwise we may either consider the complex conjugate of (1.7) or apply the transformation $y \rightarrow -y$. Hence, we can rescale t , x , and y by applying

$$t \rightarrow (\kappa J)^{2/3} t, \quad (x, y) \rightarrow (\kappa J)^{1/3} (x, y), \quad (1.8)$$

yielding

$$\begin{cases} \partial_t u = -(\mathcal{A}_{0,c} - \lambda)u & \text{in } (0, T) \times \mathbb{R}_+^2, \\ u = 0 & \text{on } \partial\mathbb{R}_+^2, \end{cases} \quad (1.9)$$

where $\mathcal{A}_{0,c}$ is the differential operator defined by

$$\mathcal{A}_{0,c} = D_x^2 + \left(D_y - \frac{1}{2}x^2\right)^2 + icy, \quad (1.10)$$

with

$$D_x = -i\partial_x, \quad D_y = -i\partial_y, \quad c \in \mathbb{R}_+,$$

and

$$c = \frac{\kappa^2}{\sigma}, \quad \lambda = \lambda_0 \equiv \frac{\kappa^{4/3}}{J^{2/3}}, \quad u(t, x, y) = \psi((\kappa J)^{-2/3}t, (\kappa J)^{-1/3}x, (\kappa J)^{-1/3}y).$$

As the operator $\mathcal{A}_{0,c}$ is defined on smooth functions only, it is necessary to extend its domain to include also weakly differentiable functions. We use the Lax-Milgram theorem to this end. Consider the sesquilinear form defined on $\tilde{V} \times \tilde{V}$ by

$$(u, v) \longmapsto \tilde{a}(u, v) = \langle D_x u, D_x v \rangle + \langle (D_y - \frac{x^2}{2})u, (D_y - \frac{x^2}{2})v \rangle + ic \int_0^{+\infty} y u \bar{v} dx dy, \quad (1.11)$$

where

$$\tilde{V} = H_0^{1,\text{mag}}(\mathbb{R}_+^2, \mathbb{C}) \cap L^2(\mathbb{R}_+^2, \mathbb{C}; y dx dy), \quad (1.12)$$

and $H_0^{1,\text{mag}}(\mathbb{R}_+^2, \mathbb{C})$ is the closure of $C_c^\infty(\mathbb{R}_+^2, \mathbb{C})$ under the norm

$$u \longmapsto \sqrt{\|u\|^2 + \|D_x u\|^2 + \|(D_y - \frac{x^2}{2})u\|^2}.$$

Here and thereafter we use $L^2(\mathbb{R}_+^2)$ to denote the usual L^2 space of real-valued functions, $L^2(\mathbb{R}_+^2, \mathbb{C})$ to denote the L^2 space of the complex-valued functions, and use $L^2(\mathbb{R}_+^2; y dx dy)$ and $L^2(\mathbb{R}_+^2, \mathbb{C}; y dx dy)$ to denote the related spaces with measure $y dx dy$. We use $\|\cdot\|$ and $\langle \cdot, \cdot \rangle$ to denote the L^2 norm and L^2 inner product on \mathbb{R}_+^2 :

$$\|u\| = \|u\|_{L^2(\mathbb{R}_+^2)} = \left(\int_{\mathbb{R}_+^2} |u|^2 dx \right)^{1/2}, \quad \langle u, v \rangle = \int_{\mathbb{R}_+^2} u \bar{v} dx.$$

Denote further by \tilde{q} the associated form

$$\tilde{V} \ni u \longmapsto \tilde{q}(u) = \tilde{a}(u, u).$$

Observing that, when $c \neq 0$, there exists $\alpha > 0$ such that

$$\tilde{q}(u) \geq \alpha \|u\|_{\tilde{V}}^2, \quad \text{for all } u \in \tilde{V},$$

we use the Lax-Milgram theorem to extend $\mathcal{A}_{0,c}$ into the unbounded operator \mathcal{A}_c^+ on $L^2(\mathbb{R}_+^2, \mathbb{C})$ whose domain $D(\mathcal{A}_c^+)$ is the set of the $u \in \tilde{V}$ such that the map

$$C_c^\infty(\mathbb{R}_+^2, \mathbb{C}) \ni v \longmapsto \tilde{a}(u, v)$$

extends as an anti-linear form on $L^2(\mathbb{R}_+^2, \mathbb{C})$, which we denote by $\langle \mathcal{A}_c^+ u, \cdot \rangle$. Hence we have

$$D(\mathcal{A}_c^+) = \{u \in \tilde{V} : \mathcal{A}_c^+ u \in L^2(\mathbb{R}_+^2, \mathbb{C})\},$$

and

$$\tilde{a}(u, v) = \langle \mathcal{A}_c^+ u, v \rangle, \quad \forall u \in D(\mathcal{A}_c^+), \quad \forall v \in \tilde{V}.$$

Once the definition of the extended operator \mathcal{A}_c^+ has been formulated, we may write

$$\mathcal{A}_c^+ = D_x^2 + (D_y - \frac{1}{2}x^2)^2 + icy. \quad (1.13)$$

Note that \mathcal{A}_c^+ is not self-adjoint but we have

$$(\mathcal{A}_c^+)^* = \mathcal{A}_{-c}^+.$$

In the present contribution we analyze the spectrum of \mathcal{A}_c^+ , denoted by $\sigma(\mathcal{A}_c^+)$, and the semi-group associated with \mathcal{A}_c^+ , which we denote by $\exp(-t\mathcal{A}_c^+)$.

By the methods of [4], we can easily prove the following :

Proposition 1.1. *For any $c > 0$, \mathcal{A}_c^+ has a compact resolvent. Moreover, if $E_0(\beta)$ denotes the ground state energy of the anharmonic oscillator*

$$-\frac{d^2}{dx^2} + \left(\frac{x^2}{2} - \beta\right)^2,$$

and if

$$E_0^* = \inf_{\beta \in \mathbb{R}} E_0(\beta), \quad (1.14)$$

then

$$\sigma(\mathcal{A}_c^+) \subset \{\lambda \in \mathbb{C}, \operatorname{Re} \lambda \geq E_0^*\}. \quad (1.15)$$

As mentioned earlier, our interest is in the effect that the Dirichlet boundary condition in (1.9) has on the spectrum $\sigma(\mathcal{A}_c^+)$ and on the semigroup $\exp(-t\mathcal{A}_c^+)$. Thus, it is interesting to compare them with the analogous entities for the whole-plane problem. Recall that in [4] we denoted by \mathcal{A} the operator corresponding to the extension of $A_{0,c}$ from $C_c^\infty(\mathbb{R}^2, \mathbb{C})$ to its closure under the associated graph norm.

Recall further, that for the whole plane problem the spectrum of \mathcal{A} is invariant under translations in a direction parallel to the imaginary axis. This property, together with the fact that the resolvent of \mathcal{A} is compact, has the consequence that $\sigma(\mathcal{A})$ must be empty and that the decay of the semigroup $\exp(-t\mathcal{A})$ is faster than any exponential rate. On the other hand, for the half-plane problem discussed in the present paper, such an invariance principle doesn't hold for $\sigma(\mathcal{A}_c^+)$. Hence, we do not expect $\sigma(\mathcal{A}_c^+)$ to be empty. Instead, we expect that eigenvalues of \mathcal{A}_c^+ would exist, and that the system of corresponding eigenfunctions would be complete in $L^2(\mathbb{R}_+^2, \mathbb{C})$. We provide here a proof of the former statement in the asymptotic regime $c \rightarrow +\infty$, and leave the proof of the latter, as well as the discussion of the limit $c \rightarrow 0$, to future research.

The main results of this paper are the following:

Theorem 1.2. *There exists $c_0 \geq 0$ such that if $c \geq c_0$ then*

$$\sigma(\mathcal{A}_c^+) \neq \emptyset.$$

Furthermore, there exists $\mu(c) \in \sigma(\mathcal{A}_c^+)$ which behaves according to the following expansion as $c \rightarrow +\infty$:

$$\mu(c) \sim c^{2/3} \exp(i\frac{\pi}{3})\alpha_0 + \lambda_1 \exp(-i\frac{\pi}{6})c^{-1/3} + \mathcal{O}(c^{-5/6}), \quad (1.16)$$

where $-\alpha_0$ is the rightmost zero point of the Airy's function [1], and λ_1 is given by (4.13). Finally, let

$$\mu_m(c) = \inf_{z \in \sigma(\mathcal{A}_c^+)} \operatorname{Re} z. \quad (1.17)$$

Then, for all $c \geq c_0$ we have

$$\mu_m(c) \sim \operatorname{Re} \mu(c) + \mathcal{O}(c^{-5/6}). \quad (1.18)$$

The lower bound of (1.18) is obtained in Proposition 3.5, whereas the upper bound is proved in Corollary 6.2.

The next result is valid for all $c > 0$.

Theorem 1.3. *If $\sigma(\mathcal{A}_c^+) \neq \emptyset$, then*

$$\lim_{t \rightarrow +\infty} -\frac{\log \|\exp(-t\mathcal{A}_c^+)\|}{t} = \mu_m(c). \quad (1.19)$$

One can easily obtain from (1.16) and (1.18) the critical current J_c , for which the normal state loses its stability in the case when the electric current is perpendicular to the surface and the applied magnetic field is parallel to it. In fact for $0 < \lambda < \mu_m$ the zero solution of (1.9) is stable, whereas for $\lambda > \mu_m$ it is unstable, as is manifested by Proposition 7.1 (see also [3] for the discussion of the case where the magnetic field is absent). From (1.16) and (1.18) we obtain that

$$\frac{J_c}{\sigma} = c\mu_m^{-3/2} = \left[\frac{\alpha_0}{2} + \frac{\sqrt{3}\lambda_1}{2c} + \mathcal{O}(c^{-3/2}) \right]^{-3/2}. \quad (1.20)$$

Note that the critical current is independent of the applied magnetic field $\mathcal{H}^e = h\mathbf{i}_z$, whose sole effect is to translate the x coordinate by h/J .

From (1.20) we see that for a large value of c

$$\frac{J_c}{\sigma} \sim \left[\frac{\alpha_0}{2} \right]^{-3/2} \left[1 - \frac{3\sqrt{3}\lambda_1}{2\alpha_0 c} + \mathcal{O}(c^{-3/2}) \right]. \quad (1.21)$$

When neglecting the induced magnetic field the critical current can be evaluated for all $c > 0$ (cf. [14, 3]). In this case we have

$$\frac{J_c}{\sigma} = \left[\frac{\alpha_0}{2} \right]^{-3/2}.$$

Consequently, one can persuasively argue that the simplified model, where the magnetic field is neglected [14, 3, 18, 17, 19], can be obtained from the linearized Ginzburg-Landau system in the limit $c \rightarrow +\infty$.

The above results significantly differ from those obtained previously for the problem in the entire plane [4]. In the case of the half-plane the critical current is positive, meaning that the normal state becomes unstable for sufficiently small currents. In contrast, for the entire plane problem the normal state is stable for every non-zero current. This suggests that for a sample of finite size, if the current is lowered below some critical value, then instability would be initiated near the boundary. For the reduced model [14], where the magnetic field is neglected, it has been demonstrated numerically for a one-dimensional setting, that the emerging superconducting phase eventually captures the whole domain. The presence of a magnetic field, may however have a stabilizing effect, and hence, one can expect a stable bifurcation of a thin surface layer of superconducting material from the normal state.

In this paper we focus our attention on the Dirichlet boundary condition $\psi = 0$ on $\partial\mathbb{R}_+^2$ given in (1.1) as it represents a normal/superconducting interface. Other boundary conditions such as the Neumann condition $\frac{\partial\psi}{\partial\nu} = 0$ on $\partial\mathbb{R}_+^2$, or the generalized Neumann condition $(\nabla - i\mathbf{A})\psi \cdot \nu = 0$, are also of great interest to mathematicians and physicists.

As a final remark, note that the present contribution focuses on the special case where the electric current is perpendicular to the surface and the applied magnetic field is parallel to it. Though this case is encountered frequently in experiments, where the potential is often constant on the surface, the more general case where the electric current and the applied magnetic field are arbitrarily directed is also of significant interest.

The rest of this work is arranged as follows: In the next section we transform \mathcal{A}_c^+ into a new operator via analytic dilation and prove the relation between $\sigma(\mathcal{A}_c^+)$ and the spectrum of the transformed operator. In § 3 we outline the scheme of the proof of Theorem 1.2. In § 4, we use formal asymptotics to estimate the principal eigenvalue of \mathcal{A}_c^+ and the corresponding eigenmode (the approximate eigenmode is usually called quasimode in the literature). To bound the error resulting from the approximation of the exact eigenmode by the quasimode, we prove in § 5 some elliptic estimates. These estimates are used in § 6 to complete the proof of Theorem 1.2. Finally, in § 7, we prove Theorem 1.3.

Acknowledgements. Y. Almog was supported by NSF grant DMS 0604467. The work of X. B. Pan was partially supported by the National Natural Science Foundation of China grant no. 10871071, and the National Basic Research Program of China grant no. 2006CB805902.

2. ANALYTIC DILATION

In this section, we assume $c \neq 0$ and apply a technique inspired by the wisdom acquired during the study of the resonances of some Schrödinger operators by Aguilar and Combes [2] or Combes and Thomas [7] (see also Simon [20, p. 498-500] for a survey and Davies [8, Lemma 5]).

For technical reasons it is easier to consider, instead of \mathcal{A}_c^+ , the unitarily equivalent operator:

$$\mathcal{P} = e^{-ix^2y/2} \mathcal{A}_c^+ e^{ix^2y/2} = (D_x + xy)^2 + D_y^2 + icy. \quad (2.1)$$

Its associated sesquilinear form is given by

$$a(u, v) = \langle (D_x + yx)u, (D_x + yx)v \rangle + \langle D_y u, D_y v \rangle + ic \langle y^{1/2}u, y^{1/2}v \rangle,$$

whose domain is given by

$$V = \{u \in L^2(\mathbb{R}_+^2, \mathbb{C}) : (D_x + yx)u \in L^2(\mathbb{R}_+^2, \mathbb{C}), \\ D_y u \in L^2(\mathbb{R}_+^2, \mathbb{C}), y^{1/2}u \in L^2(\mathbb{R}_+^2, \mathbb{C}), u(\cdot, 0) = 0\}.$$

The condition $u(\cdot, 0) = 0$ is to be understood in the sense of trace. Note that $V \subset H_{\text{loc}}^1(\overline{\mathbb{R}_+^2}, \mathbb{C})$, hence every $u \in V$ has trace lying in $H_{\text{loc}}^{1/2}(\partial\mathbb{R}_+^2, \mathbb{C})$. For later reference we note the obvious fact that V is dense in $L^2(\mathbb{R}_+^2, \mathbb{C})$.

Let $\theta \in \mathbb{C}$. Using a variant of the argument in [2], we introduce the dilation operator

$$u \mapsto (U(\theta)u)(x, y) = e^{-\theta/2} u(e^\theta x, e^{-2\theta} y). \quad (2.2)$$

Set then

$$\mathcal{P}_\theta := U(\theta)^{-1} \mathcal{P} U(\theta) = e^{2\theta} (D_x + yx)^2 - e^{-4\theta} \partial_y^2 + ic e^{2\theta} y, \quad (2.3)$$

with the associated sesquilinear form a_θ given by

$$(u, v) \mapsto a_\theta(u, v) = e^{2\theta} \langle (D_x + yx)u, (D_x + yx)v \rangle \\ + e^{-4\theta} \langle D_y u, D_y v \rangle + ic e^{2\theta} \langle y^{1/2}u, y^{1/2}v \rangle. \quad (2.4)$$

The domain of all sesquilinear forms a_θ is still given by $V \times V$ independently of θ . We denote by q_θ the corresponding quadratic form :

$$q_\theta(u) = a_\theta(u, u).$$

Lemma 2.1. *If θ varies in \mathbb{R} , then the spectrum of \mathcal{P}_θ is independent of θ .*

Proof. Since $U(\theta)$ is unitary for all $\theta \in \mathbb{R}$, it follows from (2.3) that \mathcal{P}_θ is unitarily equivalent to \mathcal{P} for all $\theta \in \mathbb{R}$. It follows that the spectrum of \mathcal{P}_θ is independent of θ if we restrict θ to the real line. ■

However, as we need to consider complex values of θ in the sequel, we have to first establish holomorphic dependence on $\theta \in \mathbb{C}$, in a sense to be made more precise (cf. [15]), of $\sigma(\mathcal{P}_\theta)$. To this end we first prove the following lemma.

Lemma 2.2. *Let*

$$D_0 = \{z \in \mathbb{C} : -\frac{\pi}{8} < \operatorname{Im} z < \frac{\pi}{24}\}.$$

Then, the forms

$$\tilde{q}_\theta := e^{-2\theta - i\pi/4} q_\theta, \quad \theta \in D_0, \quad (2.5)$$

constitute a holomorphic family of type (a).

Proof. For the precise definition of a type (a) holomorphic family of forms the reader is referred to [15, section VII.4.2]. To show that the forms (2.5) are of type (a), we have to show that, for all $\theta \in D_0$, the form \tilde{q}_θ given in (2.5) is sectorial and closed (see [15, p.310, 313]).

To show \tilde{q}_θ is sectorial, we need to estimate the size of the numerical range of the form q_θ , namely the set

$$\{z \in \mathbb{C} : \exists u \in V \text{ with } \|u\| = 1 \text{ and } z = q_\theta(u)\}. \quad (2.6)$$

For every $u \in V$ we have

$$q_\theta(u) = e^{2\theta} (\|(D_x + yx)u\|^2 + ic\|y^{1/2}u\|^2) + e^{-4\theta} \|\partial_y u\|^2. \quad (2.7)$$

Consequently, we obtain that

$$\min\{-4 \operatorname{Im} \theta, 2 \operatorname{Im} \theta\} = \alpha_m \leq \arg q_\theta(u) \leq \alpha_M = \max\{-4 \operatorname{Im} \theta, 2 \operatorname{Im} \theta + \frac{\pi}{2}\}, \quad (2.8)$$

and hence

$$|\arg e^{-2\theta - i\pi/4} q_\theta(u)| \leq \max\{\frac{\pi}{4}, |6 \operatorname{Im} \theta + \frac{\pi}{4}|\}.$$

Since

$$|6 \operatorname{Im} \theta + \frac{\pi}{4}| < \frac{\pi}{2}, \quad \forall \theta \in D_0,$$

we have

$$|\arg \tilde{q}_\theta(u)| = |\arg e^{-2\theta - i\pi/4} q_\theta(u)| < \frac{\pi}{2}, \quad \forall \theta \in D_0,$$

from which sectoriality easily follows.

Finally, we need to verify that the form \tilde{q}_θ is closed. This is an immediate consequence of (2.7), the completeness of the space $H_0^{1, \operatorname{mag}}(\mathbb{R}_+^2, \mathbb{C})$, and the definition of closeness for sesquilinear forms (cf. [15, Section VI.1.3]). ■

Remark 2.3. *One can show that \mathcal{P}_θ is a type B holomorphic family of operators (see [15, p.395] for the definition) in the larger region*

$$-\frac{\pi}{6} < \operatorname{Im} \theta < \frac{\pi}{12}.$$

For the purpose of this work, it suffices to prove this property for every $\theta \in D_0$, a fact which follows immediately from Lemma 2.2 and [15, Theorem VII.4.2].

Using the same technique as in the proof of [4, Proposition 2.4], the resolvent of \mathcal{P}_θ can be shown to be compact for every $\theta \in D_0$. Hence, we may use [15, Theorem VII.1.9] together with the fact, proved in Lemma 2.1, that $\sigma(\mathcal{P}_\theta)$ is independent of θ for $\theta \in \mathbb{R}$, to obtain the following:

Proposition 2.4. *The spectrum of \mathcal{P}_θ is independent of θ in D_0 .*

As a corollary we get, by taking $\theta = -i\frac{\pi}{12}$, the following:

Corollary 2.5. *The spectrum of \mathcal{A}_c^+ satisfies:*

$$\sigma(\mathcal{A}_c^+) \subset \{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \geq \frac{1}{2}|c|^{2/3}\alpha_0\}, \quad (2.9)$$

where α_0 is the ground-state energy of the Dirichlet realization of the Airy operator

$$\mathcal{L} = D_y^2 + y, \quad (2.10)$$

on \mathbb{R}_+ .

We recall that $-\alpha_0$ is the rightmost zero point of the standard Airy function A_i , and

$$\psi_0(y) = A_i(y - \alpha_0) \quad (2.11)$$

is the eigenfunction of the operator \mathcal{L} on \mathbb{R}_+ with Dirichlet boundary condition at $y = 0$, associated with the eigenvalue α_0 . More generally, if $-\alpha_k \in \mathbb{R}_-$ ($k \in \mathbb{N}$) denotes the $(k+1)$ 'th zero of Airy's function, then

$$\psi_k(y) = A_i(y - \alpha_k) \quad (2.12)$$

is the eigenfunction of \mathcal{L} on \mathbb{R}_+ with Dirichlet boundary condition at $y = 0$, associated with α_k .

3. THE SCHEME OF THE PROOF OF THE MAIN RESULTS

In this section we outline the main steps of the proof of Theorem 1.2. Our goal is, thus, to obtain an estimate, in the large c limit, for the real part of the leftmost eigenvalue of \mathcal{A}_c^+ in \mathbb{C} . An immediate corollary would then be that $\sigma(\mathcal{A}_c^+) \neq \emptyset$.

Instead of dealing with $\sigma(\mathcal{A}_c^+)$, it is more convenient to analyze the spectrum of the operator \mathcal{P}_θ which is given by (2.3), and is obtained from \mathcal{A}_c^+ using analytic dilation. For

$$\theta = -i\frac{\pi}{12}.$$

we have

$$\mathcal{P}_{-i\frac{\pi}{12}} = e^{i\pi/3}(D_y^2 + cy) + e^{-i\pi/6}(D_x + xy)^2.$$

Special emphasis should be given to the fact that $\mathcal{P}_{-i\frac{\pi}{12}}$ is *not* unitarily equivalent to \mathcal{A}_c^+ . Hence, analytic dilation facilitates the analysis of the spectrum of \mathcal{A}_c^+ , but the decay of the associated semi-group has to be obtained using a different approach. Since we consider large values of c it is natural to introduce the small parameter

$$\epsilon = \frac{1}{c}.$$

We define another operator via the (real) dilation

$$\mathcal{B}_\epsilon := \epsilon^{2/3} e^{i\pi/6} U(-\ln \frac{\epsilon}{6})^{-1} \mathcal{P}_{-i\frac{\pi}{12}} U(-\ln \frac{\epsilon}{6}), \quad (3.1)$$

where U is defined in (2.2). Explicitly we have

$$\mathcal{B}_\epsilon := \epsilon(D_x + xy)^2 + i(D_y^2 + y). \quad (3.2)$$

The domain of \mathcal{B}_ϵ is

$$D(\mathcal{B}_\epsilon) = \{u \in \tilde{V} : \mathcal{B}_\epsilon u \in L^2(\mathbb{R}_+^2, \mathbb{C})\},$$

where \tilde{V} is given by (1.12).

From the foregoing discussion it follows that $\sigma(\mathcal{A}_c^+)$ can easily be obtained from $\sigma(\mathcal{B}_\epsilon)$. In fact, any property we prove for the latter, including the next proposition, can be translated into a similar property of the former.

Proposition 3.1. *There exists $\epsilon_0 > 0$ such that for $\epsilon \in (0, \epsilon_0)$ we have $\sigma(\mathcal{B}_\epsilon) \neq \emptyset$. Furthermore, there exists $\lambda \in \sigma(\mathcal{B}_\epsilon)$ such that*

$$|\lambda - i\alpha_0 - \epsilon\lambda_1| < C\epsilon^{3/2} \quad \text{for all } 0 < \epsilon < \epsilon_0, \quad (3.3)$$

where $-\alpha_0$ is the rightmost zero point of the Airy's function [1], and λ_1 is given by (4.13) in Section 4.

The proof of the proposition is divided into four main steps.

Step 1.

The first step entails the construction of quasimodes. Note that for a self-adjoint operator, a quasimode provides us with an upper bound for the bottom of the spectrum, via a variational principle, and hence the spectrum cannot be empty. For the non-self-adjoint operator \mathcal{B}_ϵ , we construct the quasimodes in the next section, where we also obtain a formal asymptotic expansion, presented in Proposition 3.2, for the associated eigenvalue.

Denote by $\mathcal{S}(\overline{\mathbb{R}_+^2})$ the space of the fast decay smooth functions defined on \mathbb{R}_+^2 (cf. [12] for the precise definition), and by $\mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$ the corresponding space of complex-valued functions.

Proposition 3.2. *There exist u_j ($j = 0, 1$) in $\mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C}) \cap D(\mathcal{B}_\epsilon)$ with $\|u_0\| = 1$ and $\lambda_1 \in \mathbb{R}$, such that for*

$$\begin{cases} u_\epsilon^{\text{app}}(x, y) &= u_0(x, y) + \epsilon u_1(x, y), \\ \gamma_\epsilon &= i\alpha_0 + \epsilon\lambda_1, \end{cases} \quad (3.4)$$

we have

$$(\mathcal{B}_\epsilon - \gamma_\epsilon) u_\epsilon^{\text{app}} = \epsilon^2 f_\epsilon, \quad (3.5)$$

with f_ϵ being uniformly bounded in $(0, \epsilon_0)$ for any norm on $\mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$.

Proposition 3.2 is proved in Section 4.

Remark 3.3. *To describe the topology in $\mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$, it is sufficient to use the family of norms*

$$f \longmapsto p_k(f) = \sum_{p+q+r+s \leq k} \|x^p y^q \partial_x^r \partial_y^s f\|, \quad (3.6)$$

for $k \in \mathbb{N}$. We use this definition in the sequel whenever a norm appears in our calculations, except for the rare cases where the exact form of the norm should be introduced.

Step 2.

We next show the existence of $\rho > 0$ such that the circle

$$\mathcal{C}_{\epsilon, \rho} = \{\lambda \in \mathbb{C} : |\lambda - \gamma_\epsilon| = \rho\epsilon^{3/2}\} \quad (3.7)$$

does not intersect with the spectrum of \mathcal{B}_ϵ for sufficiently small ϵ . Once this disjointedness is established, we can define the associated projector :

$$\Pi_\epsilon = \frac{i}{2\pi} \oint_{\mathcal{C}_{\epsilon, \rho}} (\mathcal{B}_\epsilon - \lambda)^{-1} d\lambda. \quad (3.8)$$

Step 3.

Then we show that the projector Π_ϵ does not identically vanish. To this end it suffices to show that

$$\Pi_\epsilon u_\epsilon^{\text{app}} \neq 0 \quad (3.9)$$

for the same u_ϵ^{app} given in (3.4).

To prove (3.9) we represent (3.5) in the form

$$u_\epsilon^{\text{app}} = \frac{\epsilon^2}{\lambda - \gamma_\epsilon} f_\epsilon - \frac{1}{\lambda - \gamma_\epsilon} (\mathcal{B}_\epsilon - \lambda) u_\epsilon^{\text{app}},$$

which leads to

$$i(\mathcal{B}_\epsilon - \lambda)^{-1} u_\epsilon^{\text{app}} = \frac{i\epsilon^2}{\lambda - \gamma_\epsilon} (\mathcal{B}_\epsilon - \lambda)^{-1} f_\epsilon - \frac{i}{\lambda - \gamma_\epsilon} u_\epsilon^{\text{app}}.$$

Integrating the above identity with respect to λ around $\mathcal{C}_{\epsilon, \rho}$ in the positive direction we obtain

$$\Pi_\epsilon u_\epsilon^{\text{app}} = u_\epsilon^{\text{app}} + \frac{i\epsilon^2}{2\pi} \oint_{\mathcal{C}_{\epsilon, \rho}} \frac{1}{\lambda - \gamma_\epsilon} (\mathcal{B}_\epsilon - \lambda)^{-1} f_\epsilon d\lambda. \quad (3.10)$$

It then remains to show that the right-hand-side of (3.10) does not identically vanish.

Step 4.

Using (3.10) we see that to prove (3.9), it is sufficient to uniformly control the L^2 norm of $(\mathcal{B}_\epsilon - \lambda)^{-1} f_\epsilon$ for all λ on the circle $\mathcal{C}_{\epsilon, \rho}$. In fact we prove the following

Proposition 3.4. *There exist positive constants ρ , C and ϵ_0 and $k_0 \in \mathbb{N}$ such that, for all $\epsilon \in (0, \epsilon_0]$, $\lambda \in \mathcal{C}_{\epsilon, \rho}$, and $f \in \mathcal{S}(\overline{\mathbb{R}}_+^2, \mathbb{C})$ we have*

$$\|(\mathcal{B}_\epsilon - \lambda)^{-1} f\| \leq C\epsilon^{-3/2} p_{k_0}(f), \quad (3.11)$$

where p_{k_0} is the norm defined in (3.6).

Proposition 3.4 will be proved at the end of Section 4.

We can then conclude from (3.10) and (3.11) the existence of positive constants C and ϵ_0 , such that, for any $\epsilon \in (0, \epsilon_0]$,

$$\|\Pi_\epsilon u_\epsilon^{\text{app}}\| \geq 1 - C\epsilon^{1/2}. \quad (3.12)$$

Then for small ϵ one can conclude (3.9) from (3.12).

To complete steps 2 and 4, we analyze the equation :

$$(\mathcal{B}_\epsilon - \lambda)w = f, \quad (3.13)$$

with $f \in \mathcal{S}(\overline{\mathbb{R}}_+^2, \mathbb{C})$ and $w \in D(\mathcal{B}_\epsilon)$. We perform this task in Section 5, where we derive various estimates of the solution w of (3.13). In addition, we prove there the following result:

Proposition 3.5. *Let*

$$D_{b, \epsilon} = \{\lambda \in \mathbb{C} : 0 < \text{Re } \lambda \leq \epsilon\lambda_1 - b\epsilon^{3/2} \text{ and } 0 < \text{Im } \lambda - \alpha_0 \leq 3\lambda_1\epsilon\}. \quad (3.14)$$

Then, there exist positive b_0 and ϵ_0 such that for all $\epsilon \in (0, \epsilon_0)$ and $b > b_0$ we have

$$D_{b, \epsilon} \cap \sigma(\mathcal{B}_\epsilon) = \emptyset.$$

Note that since

$$\text{Im } \langle \mathcal{B}_\epsilon u, u \rangle = \langle \mathcal{L}u, u \rangle \geq \alpha_0 \|u\|^2 \quad \text{and} \quad \text{Re } \langle \mathcal{B}_\epsilon u, u \rangle \geq 0,$$

it follows that

$$\sigma(\mathcal{B}_\epsilon) \subset S_{b, \epsilon}, \quad (3.15)$$

where

$$S_{b,\epsilon} = \left\{ \lambda \in \mathbb{C} \left| \begin{array}{ll} \operatorname{Re} \lambda > \epsilon \lambda_1 - b\epsilon^{3/2} & \text{and } 0 < \operatorname{Im} \lambda \leq \alpha_0 + 3\lambda_1 \epsilon \\ \operatorname{Re} \lambda > 0 & \text{and } \operatorname{Im} \lambda \geq \alpha_0 + 3\lambda_1 \epsilon \end{array} \right. \right\}. \quad (3.16)$$

The lower bound in (1.18) now immediately follows from (3.15) and (3.16).

Proposition 3.5 is proved in Section 6.

4. CONSTRUCTIONS OF QUASIMODES

The main result in this section is the following Proposition 4.1, which implies Proposition 3.2 in section 3.

Proposition 4.1. *Let \mathcal{B}_ϵ be the operator defined in (3.2). There exists a sequence*

$$\{u_j(x, y)\}_{j=0}^{+\infty} \subset \mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C}) \cap D(\mathcal{B}_\epsilon) \quad (4.1)$$

with $\|u_0\|_{L^2} = 1$ and a sequence

$$\{\lambda_j\}_{j=0}^{+\infty} \subset \mathbb{C}$$

with $i^{j+1}\lambda_j$ being real for all $j \geq 0$ and

$$\lambda_0 = i\alpha_0, \quad (4.2)$$

where α_0 is the lowest eigenvalue of the Airy operator \mathcal{L} defined in (2.10) in \mathbb{R}_+ with homogeneous Dirichlet condition at $y = 0$, such that

$$\mathcal{B}_\epsilon \left(\sum_{j=0}^{+\infty} \epsilon^j u_j(x, y) \right) \sim \left(\sum_{j=0}^{+\infty} \epsilon^j \lambda_j \right) \left(\sum_{j=0}^{+\infty} \epsilon^j u_j(x, y) \right) \quad (4.3)$$

in the sense of formal expansions in ϵ .

Remark 4.2. *We note that if ϵ had been purely imaginary, then (4.3) would have implied by the spectral theorem the existence of an eigenvalue $\lambda(\epsilon)$ admitting the complete asymptotic expansion*

$$\lambda(\epsilon) \sim \sum_{j=0}^{+\infty} \epsilon^j \lambda_j.$$

Since our interest is in the non-self-adjoint case where ϵ is real, a more complicated approach has to be adopted in order to prove the existence of an eigenvalue of \mathcal{B}_ϵ .

Proof. It is sufficient to show how one can obtain u_j and λ_j for $j \leq 2$. Higher order terms can similarly be obtained. We first derive the $\mathcal{O}(\epsilon^j)$ balance from (4.3) for every $j \geq 0$. Suppose that the sequence in (4.1) satisfies (4.3) in the sense of formal expansions in ϵ . Balancing the coefficients of ϵ^j in each side of (4.3) we obtain

$$\begin{aligned} i(D_y^2 + y)u_0 &= \lambda_0 u_0, \\ i(D_y^2 + y)u_j + (D_x + xy)^2 u_{j-1} &= \sum_{\ell+m=j} \lambda_\ell u_m, \quad j \geq 1. \end{aligned} \quad (4.4)$$

Step 1. The $\mathcal{O}(\epsilon^0)$ balance gives :

$$(i(D_y^2 + y) - \lambda_0) u_0(x, y) = 0. \quad (4.5)$$

It follows immediately that $\lambda_0 = i\alpha_0$, in accordance with (4.2). Furthermore, it follows that u_0 must be in the form

$$u_0(x, y) = \phi_0(x)\psi_0(y), \quad (4.6)$$

where $\psi_0(y)$ is the L^2 -normalized (real-valued) first eigenfunction of the Airy operator $\mathcal{L} = D_y^2 + y$, defined in (2.10), in \mathbb{R}_+ with Dirichlet boundary at $y = 0$, associated with the lowest eigenvalue α_0 . It is well-known that ψ_0 is in $\mathcal{S}(\overline{\mathbb{R}_+})$ (see [1]). We leave ϕ_0 undetermined as it cannot be obtained from (4.5).

Step 2. The $\mathcal{O}(\epsilon)$ balance reads :

$$i \left((D_y^2 + y) - \alpha_0 \right) u_1(x, y) + (D_x + xy)^2 u_0(x, y) = \lambda_1 u_0(x, y). \quad (4.7)$$

A necessary condition for the solvability of (4.7) is obtained by taking the scalar product (in the y -variable) of both sides of (4.7) by ψ_0 . Since $u_1 \in \mathcal{S}(\overline{\mathbb{R}_+}, \mathbb{C}) \cap D(\mathcal{B}_\epsilon)$ we have that $u_1(x, 0) = 0$. Integration by parts then yields

$$\int_0^{+\infty} [(D_y^2 + y - \alpha_0)u_1] \bar{\psi}_0 dy = 0.$$

Since $\|\psi_0\|_{L^2(\mathbb{R}_+)} = 1$, we have that

$$\int_0^{+\infty} [(D_x + xy)^2 \phi_0(x)] \psi_0(y)^2 dy = \mathcal{K}_0 \phi_0(x), \quad (4.8)$$

where \mathcal{K}_0 is the following differential operator

$$\mathcal{K}_0(x, D_x) = (D_x + \alpha x)^2 + (\beta^2 - \alpha^2)x^2, \quad x \in \mathbb{R}, \quad (4.9)$$

in which

$$\alpha = \int_0^{+\infty} y \psi_0(y)^2 dy, \quad \beta^2 = \int_0^{+\infty} y^2 \psi_0(y)^2 dy. \quad (4.10)$$

Combining (4.7), (4.8), and (4.9) gives

$$\mathcal{K}_0 \phi_0(x) = \lambda_1 \phi_0(x). \quad (4.11)$$

Since $y\psi_0(y)$ is not a multiple of ψ_0 , it follows from Cauchy-Schwarz inequality, that the following strict inequality holds

$$\alpha < \beta. \quad (4.12)$$

Hence \mathcal{K}_0 is an harmonic oscillator, whose lowest eigenvalue is $\sqrt{\beta^2 - \alpha^2}$. Consequently,

$$\lambda_1 = \sqrt{\beta^2 - \alpha^2}, \quad (4.13)$$

and ϕ_0 is the corresponding L^2 -normalized eigenfunction. Explicitly, ϕ_0 is a complex Gaussian :

$$\phi_0(x) = a \exp\left(\frac{i\alpha}{2}x^2\right) \exp\left(-\frac{1}{2}\sqrt{\beta^2 - \alpha^2}x^2\right), \quad (4.14)$$

where the constant a is chosen so that $\|\phi_0\|_{L^2(\mathbb{R})} = 1$.

The solvability of (4.7) now easily follows: since \mathcal{L} is self-adjoint, and all its eigenvalues are simple, it follows that

$$(D_x + xy)^2 u_0(x, y) - \lambda_1 u_0(x, y) \perp \text{Ker}\{\mathcal{L} - \alpha_0\}, \quad \forall x \in \mathbb{R}.$$

As $\mathcal{L} - \alpha_0$ is a Fredholm operator with index zero, the solvability of (4.7) readily follows. Consequently, u_1 is expressible in the form

$$u_1(x, y) = u_1^0(x, y) + \phi_1(x)\psi_0(y), \quad (4.15)$$

where u_1^0 is the unique solution of (4.7) belonging to $(\text{span}\{\psi_0\})^\perp$ for all $x \in \mathbb{R}$, i.e.,

$$\int_0^{+\infty} u_1^0(x, y)\psi_0(y)dy = 0. \quad (4.16)$$

Step 3. Consider next the $\mathcal{O}(\epsilon^2)$ balance:

$$i((D_y^2 + y) - \alpha_0)u_2(x, y) + (D_x + xy)^2u_1(x, y) = \lambda_1u_1(x, y) + \lambda_2u_0(x, y). \quad (4.17)$$

A first necessary condition for solving (4.17) is obtained by taking the scalar product in the y -variable of the both sides of (4.17) with ψ_0 . Using (4.8) and (4.15) we obtain

$$\int_0^{+\infty} [(D_x + xy)^2u_1(x, y)]\psi_0(y)dy = \mathcal{K}_0\phi_1(x) + \int_0^{+\infty} [(D_x + xy)^2u_1^0(x, y)]\psi_0(y)dy.$$

Combining the above with (4.16) then yields

$$(\mathcal{K}_0 - \lambda_1)\phi_1 = \lambda_2\phi_0 - \langle \psi_0, ((D_x + xy)^2 - \lambda_1)u_1^0(x, y) \rangle_{L^2(\mathbb{R}_+)}. \quad (4.18)$$

A second solvability condition is obtained by taking the L^2 scalar product of both sides of (4.18) with ϕ_0 in the x -variable. Using (4.11) we arrive at

$$\int_{\mathbb{R}} [(\mathcal{K}_0 - \lambda_1)\phi_1]\phi_0dx = \int_{\mathbb{R}} [(\mathcal{K}_0 - \lambda_1)\phi_0]\phi_1dx = 0.$$

Hence,

$$\begin{aligned} \lambda_2 &= \int_{\mathbb{R}} \langle \psi_0, ((D_x + xy)^2 - \lambda_1)u_1^0(x, y) \rangle_{L^2(\mathbb{R}_+)} \phi_0 dx \\ &= \langle u_0, ((D_x + xy)^2 - \lambda_1)u_1^0(x, y) \rangle_{L^2(\mathbb{R}_+^2)}. \end{aligned} \quad (4.19)$$

With this choice of λ_2 , (4.18) satisfies the necessary condition for solvability, which is also a sufficient condition. We can then choose ϕ_1 to be the unique solution of (4.18) which is orthogonal to ϕ_0 .

Substituting ϕ_1 into (4.15) we obtain u_1 . Then, (4.17) satisfies the necessary condition for solvability, which is also sufficient by the same argument applied in Step 2. Thus, the solution of (4.17) assumes the form

$$u_2(x, y) = u_2^0(x, y) + \phi_2(x)\psi_0(y), \quad (4.20)$$

where u_2^0 is uniquely determined by the orthogonality condition

$$\int_0^{+\infty} u_2^0(x, y)\psi_0(y)dy = 0. \quad (4.21)$$

It is easy to show that the u_j 's are all in $\mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$ in view of the fact that both \mathcal{L} and \mathcal{K}_0 include an unbounded potential as y and $|x|$ respectively tend to infinity.

This completes the proof of Proposition 4.1. ■

Proof of Proposition 3.2.

Proposition 3.2 readily follows from Proposition 4.1. ■

5. RESOLVENT ESTIMATES

Our goal in this section is to prove Proposition 3.4 by estimating the resolvent of \mathcal{B}_ϵ for $\lambda \in D_{b,\epsilon}$, where $D_{b,\epsilon} \subset \mathbb{C}$ was given by (3.14).

5.1. Preliminary estimates.

We begin by establishing the following approximation.

Lemma 5.1. *Let ϕ be a real-valued C^∞ function which together with its derivatives is bounded for $x \in \mathbb{R}$, and $w \in \mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C}) \cap D(\mathcal{B}_\epsilon)$. Then*

$$\int_{\mathbb{R}_+^2} \phi(x)^2 (\mathcal{B}_\epsilon w) \bar{w} \, dx dy = \|\phi D_y w\|^2 + \|y^{1/2} \phi w\|^2 + \epsilon W, \quad (5.1)$$

where $W \in L^2(\mathbb{R}_+^2, \mathbb{C})$ satisfies

$$\|W\| \leq \|(D_x + xy)(\phi w)\|^2 + \|\phi' w\|^2. \quad (5.2)$$

Proof. Since w vanishes on $\partial\mathbb{R}_+^2$, we can integrate by parts and the term involving the integral on the boundary $\partial\mathbb{R}_+^2$ vanishes. We therefore have

$$\begin{aligned} \int_{\mathbb{R}_+^2} \phi(x)^2 (\mathcal{B}_\epsilon w) \bar{w} \, dx dy &= \epsilon \int_{\mathbb{R}_+^2} \{|\phi(D_x + xy)w|^2 + 2i\phi\phi' \bar{w}(D_x + xy)w\} \, dx dy \\ &\quad + i \int_{\mathbb{R}_+^2} (|\phi D_y w|^2 + y|\phi w|^2) \, dx dy. \end{aligned} \quad (5.3)$$

Taking the imaginary part of (5.3) we obtain :

$$\begin{aligned} &\operatorname{Im} \int_{\mathbb{R}_+^2} \phi(x)^2 (\mathcal{B}_\epsilon w) \bar{w} \, dx dy \\ &= \|\phi D_y w\|^2 + \|y^{1/2} \phi w\|^2 + 2\epsilon \operatorname{Re} \int_{\mathbb{R}_+^2} \phi\phi' [\bar{w}(D_x + xy)w] \, dx dy. \end{aligned} \quad (5.4)$$

The integral in the last term in the right side can be rewritten as :

$$\operatorname{Re} \int_{\mathbb{R}_+^2} \phi\phi' [\bar{w}(D_x + xy)w] \, dx dy = \operatorname{Re} \int_{\mathbb{R}_+^2} \phi' [\bar{w}(D_x + xy)(\phi w)] \, dx dy.$$

Consequently, we may conclude that

$$\left| \operatorname{Re} \int_{\mathbb{R}_+^2} \phi\phi' [\bar{w}(D_x + xy)w] \, dx dy \right| \leq \|(D_x + xy)(\phi w)\| \|\phi' w\|. \quad (5.5)$$

For the real part of the integral on the left-hand-side of (5.1) we have

$$\begin{aligned} &\operatorname{Re} \int \phi(x)^2 (\mathcal{B}_\epsilon w) \bar{w} \, dx dy \\ &= \epsilon \int_{\mathbb{R}_+^2} \{|\phi(D_x + xy)w|^2 + i\phi\phi' [\bar{w}(D_x + xy)w - w\overline{(D_x + xy)w}]\} \, dx dy \\ &= \epsilon \|\phi(D_x + xy)w - i\phi' w\|^2 - \epsilon \|\phi' w\|^2 = \epsilon \|(D_x + xy)(\phi w)\|^2 - \epsilon \|\phi' w\|^2. \end{aligned} \quad (5.6)$$

Combining (5.5) and (5.6) yields

$$\left| \int_{\mathbb{R}_+^2} \phi(x)^2 (\mathcal{B}_\epsilon w) \bar{w} \, dx dy - \|\phi D_y w\|^2 + \|y^{1/2} \phi w\|^2 \right| \leq 2\epsilon (\|(D_x + xy)(\phi w)\|^2 + \|\phi' w\|^2), \quad (5.7)$$

which completes the proof of the lemma. ■

We continue by establishing the following estimates, which are valid for λ values in a much greater range than those proved in the next subsection.

Lemma 5.2. *Let $(f, w) \in S(\overline{\mathbb{R}_+^2}, \mathbb{C}) \times D(\mathcal{B}_\epsilon)$ satisfy (3.13). Given any $C > 0$, there exist constants ϵ_0 and \widehat{C} , and, for any half-integer $k \geq 1/2$, a constant C_k , such that, for any*

$$\lambda = i\alpha_0 + \epsilon\mu_1 \quad (5.8)$$

with

$$|\mu_1| \leq C, \quad \epsilon \in (0, \epsilon_0],$$

we have

$$|x|^k w \in L^2(\mathbb{R}_+^2, \mathbb{C}), \quad |y|^k w \in L^2(\mathbb{R}_+^2, \mathbb{C}), \quad x\partial_x w \in L^2(\mathbb{R}_+^2, \mathbb{C}),$$

with

$$\| |x|^k w \| \leq C_k \left[\|w\| + \epsilon^{-1} \|(|x|^{k-1/2} + 1)f\| \right], \quad (5.9a)$$

$$\| |y|^k w \| \leq C_k \left[\|w\| + \|(y^{k-1/2} + 1)f\| \right], \quad (5.9b)$$

$$\| |x|^k \partial_x w \| \leq C_k \left[\|w\| + \epsilon^{-1} \|(|x|^{k+3/2} + y^{k+3/2} + 1)f\| \right], \quad (5.9c)$$

$$\| |y|^k \partial_x w \| \leq C_k \left[\|w\| + \epsilon^{-1} \|(|x|^{5/2} + y^{k+1} + 1)f\| + \| |y|^{k-1/2} \partial_x f \| \right], \quad (5.9d)$$

$$\| \partial_{xx} w \| \leq \widehat{C} \left[\|w\| + \epsilon^{-1} \|(|x|^{7/2} + y^{7/2} + 1)f\| + \| |y|^{3/2} \partial_x f \| + \epsilon^{-1} \| \partial_x f \| \right]. \quad (5.9e)$$

Proof. Step 1. We first obtain an estimate of $\|(\nabla + ixy\hat{i}_y)w\|$.

To this end, let $v \in \tilde{V}$ be compactly supported in the interior of \mathbb{R}_+^2 , where \tilde{V} was defined in (1.12). Multiplying (3.13) by \bar{v} and integrating by parts yields

$$\int_{\mathbb{R}_+^2} \{ \epsilon(D_x + xy)w \cdot \overline{(D_x + xy)v} + iD_y w \cdot \overline{D_y v} + iyw\bar{v} - \lambda w\bar{v} \} dx dy = \int_{\mathbb{R}_+^2} f\bar{v} dx dy. \quad (5.10)$$

For $n \geq 1$, let $\eta_n \in C^\infty(\mathbb{R}, [0, 1])$ satisfy

$$\eta_n(t) = \begin{cases} 0 & \text{if } |t| < 1, \\ 1 & \text{if } 2 < |t| < 4n, \\ 0 & \text{if } 6n < |t|, \end{cases} \quad (5.11)$$

and

$$|\eta_n'| \leq \frac{C_0}{n}.$$

Let $m \geq 0$ and set $v(x, y) = |x|^m \eta_n^2(x) w(x, y)$ in (5.10). We compute

$$\begin{aligned} (D_x + xy)w \cdot \overline{(D_x + xy)v} &= |(D_x + xy)(|x|^{m/2} \eta_n w)|^2 - |w(|x|^{m/2} \eta_n)'|^2 \\ &\quad + 2i|x|^{m/2} \eta_n (|x|^{m/2} \eta_n)' \operatorname{Re} [\bar{w}(D_x + xy)w]. \end{aligned}$$

We can thus write (5.10) in the form

$$\begin{aligned} &\epsilon \int_{\mathbb{R}_+^2} \{ |(D_x + xy)(|x|^{m/2} \eta_n w)|^2 - |w(|x|^{m/2} \eta_n)'|^2 \} dx dy \\ &+ i \int_{\mathbb{R}_+^2} \{ 2\epsilon |x|^{m/2} \eta_n (|x|^{m/2} \eta_n)' \operatorname{Re} [\bar{w}(D_x + xy)w] + |x|^m \eta_n^2 |D_y w|^2 + y |x|^m \eta_n^2 |w|^2 \} dx dy \quad (5.12) \\ &- (i\alpha_0 + \epsilon\mu_1) \int_{\mathbb{R}_+^2} |x|^m \eta_n^2 |w|^2 dx dy = \int_{\mathbb{R}_+^2} |x|^m \eta_n^2 f \bar{w} dx dy. \end{aligned}$$

Here we have used the assumption (5.8).

Taking the real part of (5.12) we get

$$\begin{aligned} & \epsilon \int_{\mathbb{R}_+^2} \{ |(D_x + xy)(|x|^{m/2}\eta_n w)|^2 - |w(|x|^{m/2}\eta_n)'|^2 - \operatorname{Re} \mu_1 |x|^m \eta_n^2 |w|^2 \} dx dy \\ &= \operatorname{Re} \int_{\mathbb{R}_+^2} |x|^m \eta_n^2 f \bar{w} dx dy. \end{aligned}$$

Hence we have

$$\begin{aligned} & \int_{\mathbb{R}_+^2} \left| (D_x + xy)(|x|^{m/2}\eta_n(x)w) \right|^2 dx dy \\ &= \operatorname{Re} \mu_1 \| |x|^{m/2}\eta_n w \|^2 + \| (|x|^{m/2}\eta_n)' w \|^2 + \epsilon^{-1} \operatorname{Re} \langle |x|^{m/2}\eta_n w, |x|^{m/2}\eta_n f \rangle. \end{aligned} \quad (5.13)$$

Taking the imaginary part of (5.12) yields

$$\begin{aligned} & \int_{\mathbb{R}_+^2} \{ 2\epsilon |x|^{m/2}\eta_n (|x|^{m/2}\eta_n)' \operatorname{Re} [\bar{w}(D_x + xy)w] + |x|^m \eta_n^2 |D_y w|^2 + y |x|^m \eta_n^2 |w|^2 \} dx dy \\ & - (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \int_{\mathbb{R}_+^2} |x|^m \eta_n^2 |w|^2 dx dy = \operatorname{Im} \int_{\mathbb{R}_+^2} |x|^m \eta_n^2 f \bar{w} dx dy. \end{aligned}$$

Hence,

$$\begin{aligned} & \| |x|^{m/2}\eta_n D_y w \|^2 + \| y^{1/2} |x|^{m/2}\eta_n w \|^2 \\ &= (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \| |x|^{m/2}\eta_n w \|^2 + \operatorname{Im} \langle \eta_n |x|^{m/2} w, \eta_n |x|^{m/2} f \rangle \\ & - 2\epsilon \int_{\mathbb{R}_+^2} |x|^{m/2}\eta_n (|x|^{m/2}\eta_n)' \operatorname{Re} [\bar{w}(D_x + xy)w] dx dy. \end{aligned} \quad (5.14)$$

The last term in the right side of (5.14) can be estimated by using (5.5) with $\phi(x) = |x|^{m/2}\eta_n(x)$. Thus,

$$\left| \int_{\mathbb{R}_+^2} |x|^{m/2}\eta_n (|x|^{m/2}\eta_n)' \operatorname{Re} [\bar{w}(D_x + xy)w] dx dy \right| \leq \| (D_x + xy)(|x|^{m/2}\eta_n w) \| \| (|x|^{m/2}\eta_n)' w \|.$$

From the above and (5.14) we obtain

$$\begin{aligned} \| |x|^{m/2}\eta_n D_y w \|^2 &\leq (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \| |x|^{m/2}\eta_n w \|^2 + \operatorname{Im} \langle |x|^{m/2}\eta_n w, |x|^{m/2}\eta_n f \rangle \\ & + 2\epsilon \| (D_x + xy)(|x|^{m/2}\eta_n w) \| \| (|x|^{m/2}\eta_n)' w \|. \end{aligned} \quad (5.15)$$

The necessary estimate of $\| (\nabla + ixy\hat{i}_y)w \|$ is obtained by summing up (5.13) and (5.15).

Step 2. Next we show that there exists c_1 , such that, if $\phi \in C_c^1(\mathbb{R}_+^2, \mathbb{C})$, then

$$\| |x|^{1/2}\phi \|^2 \leq c_1 \left[\| (D_x + xy)\phi \|^2 + \| D_y \phi \|^2 + \|\phi\|^2 \right]. \quad (5.16)$$

To prove (5.16), we use the same trick as in [4] (line above (2.23)) and write, for ϕ with compact support inside \mathbb{R}_+^2 , the identity

$$\int_{\mathbb{R}_+^2} \frac{x^2}{\sqrt{1+x^2}} |\phi|^2 dx dy = -i \int_{\mathbb{R}_+^2} \frac{x}{\sqrt{1+x^2}} [D_x + xy, D_y] \phi \bar{\phi} dx dy,$$

where $[D_x + xy, D_y]$ denotes the commutator of $D_x + xy$ and D_y . Expanding the bracket and after an integration by part, we obtain

$$\int_{\mathbb{R}_+^2} \frac{x^2}{\sqrt{1+x^2}} |\phi|^2 dx dy \leq 3 \int_{\mathbb{R}_+^2} (\| (D_x + xy)\phi \|^2 + \| D_y \phi \|^2 + \|\phi\|^2) dx dy. \quad (5.17)$$

Step 3. Let $(f, w) \in S(\overline{\mathbb{R}_+^2}, \mathbb{C}) \times D(\mathcal{B}_\epsilon)$. In this step we claim the existence of a constant c_2 , independent of (f, w) , such that for any non-negative real number m , and for all $0 < \epsilon \leq 1$, we have :

$$\||x|^{(m+1)/2}\eta_n w\| \leq c_2 \left[\||x|^{m/2}\eta_n w\| + \|w\| + \epsilon^{-1} \||x|^{m/2}f\| \right]. \quad (5.18)$$

Here the cut-off function η_n is defined by (5.11).

We first note that, since $w \in \tilde{V}$, by approximation, (5.16) holds for

$$\phi(x, y) = |x|^{m/2}\eta_n(x)w(x, y)$$

with $m \geq 0$. Consequently, we obtain

$$\begin{aligned} \||x|^{(m+1)/2}\eta_n(x)w\|^2 &\leq c_1 \left[\|(D_x + xy)(|x|^{m/2}\eta_n(x)w)\|^2 \right. \\ &\quad \left. + \|D_y(|x|^{m/2}\eta_n(x)w)\|^2 + \||x|^{m/2}\eta_n(x)w\|^2 \right]. \end{aligned} \quad (5.19)$$

We use Cauchy inequality to control the term

$$2\epsilon \|(D_x + xy)(|x|^{m/2}\eta_n w)\| \|(|x|^{m/2}\eta_n)' w\|$$

on the right-hand-side of (5.15), and then substitute (5.13) and (5.15) into (5.19). We then obtain the existence of $c_3 > 0$, such that

$$\begin{aligned} \||x|^{(m+1)/2}\eta_n w\|^2 &\leq c_3 \left[(\operatorname{Re} \mu_1 + \alpha_0 + \epsilon \operatorname{Im} \mu_1 + 1) \||x|^{m/2}\eta_n w\|^2 + (1 + \epsilon) \|(|x|^{m/2}\eta_n)' w\|^2 \right. \\ &\quad \left. + \epsilon^{-1} \operatorname{Re} \langle |x|^{m/2}\eta_n w, |x|^{m/2}\eta_n f \rangle + \operatorname{Im} \langle |x|^{m/2}\eta_n w, |x|^{m/2}\eta_n f \rangle \right] \\ &\leq c_3 \left[(\operatorname{Re} \mu_1 + \alpha_0 + \epsilon \operatorname{Im} \mu_1 + 3) \||x|^{m/2}\eta_n w\|^2 + (1 + \epsilon) \|(|x|^{m/2}\eta_n)' w\|^2 \right. \\ &\quad \left. + (\epsilon^{-2} + 1) \||x|^{m/2}\eta_n f\|^2 \right]. \end{aligned} \quad (5.20)$$

Recall that $\eta_n(x) = 0$ for $|x| \leq 1$, and that

$$|\eta_n'(x)| \leq \eta_{2n}(x), \quad |x|^{m/2}\eta_n(x) \leq 1 + \eta_{2n}(x) \quad \text{for } |x| \geq 2.$$

Hence

$$\begin{aligned} \|(|x|^{m/2}\eta_n)' w\|^2 &\leq \||\frac{m}{2|x|}(|x|^{m/2}\eta_n w)\|^2 + \||x|^{m/2}\eta_n'(x)w\|^2 \\ &\leq m^2 \||x|^{m/2}\eta_n w\|^2 + \||x|^{m/2}\eta_{2n}(x)w\|^2 \\ &\leq (2m^2 + 2)(\|w\|^2 + \||x|^{m/2}\eta_{2n}(x)w\|^2), \end{aligned} \quad (5.21)$$

and

$$\||x|^{m/2}\eta_n w\|^2 \leq 2\|w\|^2 + 2\||x|^{m/2}\eta_{2n}(x)w\|^2. \quad (5.22)$$

Next, we insert back (5.21) and (5.22) into (5.20), and obtain the existence of c_4 , such that for all $0 < \epsilon \leq 1$ we have

$$\begin{aligned} \||x|^{(m+1)/2}\eta_n w\|^2 &\leq c_4(1 + m^2 + \alpha_0 + |\mu_1|)(\|w\|^2 + \||x|^{m/2}\eta_{2n}(x)w\|^2) \\ &\quad + c_4\epsilon^{-2} \||x|^{m/2}\eta_n f\|^2. \end{aligned} \quad (5.23)$$

Hence, (5.18) is proved.

Step 4. Proof of (5.9a). Setting first $m = 0$ in (5.18) we obtain

$$\||x|^{1/2}\eta_n w\| \leq c_2(\|\eta_n w\| + \|w\| + \epsilon^{-1}\|f\|) \leq c_2(2\|w\| + \epsilon^{-1}\|f\|).$$

Letting n go to $+\infty$ then yields

$$\| |x|^{1/2} w \| \leq c_2(2\|w\| + \epsilon^{-1}\|f\|). \quad (5.24)$$

Consequently, (5.9a) holds with $k = 1/2$. For a general half integer k , the proof of (5.9a) now follows by recursively applying the above procedure.

Step 5. Proof of (5.9b). Let $m \geq 0$ and insert $v(x, y) = y^m \eta_n^2(y) \bar{w}(x, y)$ into (5.10) to obtain

$$\begin{aligned} & \int_{\mathbb{R}_+^2} \{ \epsilon y^m \eta_n^2(y) |(D_x + xy)w|^2 + i y^m \eta_n^2(y) |D_y w|^2 + i (y^m \eta_n^2(y))' \bar{w} D_y w \\ & \quad + i y^{m+1} \eta_n^2(y) |w|^2 - \lambda y^m \eta_n^2(y) |w|^2 \} dx dy = \int_{\mathbb{R}_+^2} y^m \eta_n^2(y) f \bar{w} dx dy. \end{aligned}$$

From the imaginary part of the above identity and (5.8) we obtain that

$$\begin{aligned} & \int_{\mathbb{R}_+^2} \{ y^m \eta_n^2(y) |D_y w|^2 + (y^m \eta_n^2(y))' \operatorname{Re}(\bar{w} D_y w) + y^{m+1} \eta_n^2(y) |w|^2 \\ & \quad - (\alpha_0 + \epsilon \operatorname{Im} \mu_1) y^m \eta_n^2(y) |w|^2 \} dx dy = \operatorname{Im} \int_{\mathbb{R}_+^2} y^m \eta_n^2(y) f \bar{w} dx dy. \end{aligned}$$

Therefore,

$$\begin{aligned} & \| y^{m/2} \eta_n(y) D_y w \|^2 + \| y^{(m+1)/2} \eta_n w \|^2 \\ & \leq (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \| y^{m/2} \eta_n(y) w \|^2 + \| (y^{m/2} \eta_n(y))' w y^{m/2} \eta_n(y) D_y w \| \\ & \quad + | \operatorname{Im} \langle \eta_n(y) y^{m/2} w, \eta_n(y) y^{m/2} f \rangle | \\ & \leq (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \| y^{m/2} \eta_n(y) w \|^2 + \frac{1}{4} \| (y^{m/2} \eta_n(y))' w \|^2 + \| y^{m/2} \eta_n(y) D_y w \|^2 \\ & \quad + \frac{1}{2} \| y^{m/2} \eta_n(y) w \|^2 + \frac{1}{2} \| y^{m/2} \eta_n(y) f \|^2. \end{aligned}$$

Hence,

$$\begin{aligned} \| y^{(m+1)/2} \eta_n w \|^2 & \leq (\alpha_0 + \epsilon \operatorname{Im} \mu_1 + \frac{1}{2}) \| y^{m/2} \eta_n(y) w \|^2 + \frac{1}{4} \| (y^{m/2} \eta_n(y))' w \|^2 \\ & \quad + \frac{1}{2} \| y^{m/2} \eta_n(y) f \|^2. \end{aligned} \quad (5.25)$$

Setting $m = 0$ in (5.25) we obtain, for sufficiently large n , that

$$\| y^{1/2} \eta_n w \|^2 \leq (\alpha_0 + \epsilon \operatorname{Im} \mu_1 + 1) \| w \|^2 + \| f \|^2.$$

Letting $n \rightarrow +\infty$ in the above inequality yields the existence of c_5 such that :

$$\| |x|^{1/2} w \|^2 \leq c_5 (\|w\|^2 + \|f\|^2).$$

Invoking inductive arguments we get (5.9b) easily.

Step 6. Proof of (5.9c). For a half integer $k \geq 1/2$ we use (5.13) with $m = 2k$, and then let $n \rightarrow +\infty$, to obtain the existence of c_6 such that :

$$\| \partial_x (|x|^k w) \|^2 \leq c_6 (\| |x|^k w \|^2 + \| w \|^2) + \epsilon^{-1} \| |x|^k w \| \| |x|^k f \| + \| |x|^{k+1} y w \|^2.$$

Combining the above inequality with (5.9a) and (5.9b) and the following inequality (which immediately follows from Young's inequality)

$$|x|^p y^2 \leq \frac{1}{p+1} [p|x|^{p+1} + y^{2p+2}] \quad (5.26)$$

completes the proof of (5.9c).

In the interest of brevity, we drop from now on the argument involving the cut-off function η_n (including the limit $n \rightarrow +\infty$).

Step 7. Proof of (5.9d). We differentiate (3.13) with respect to x to obtain

$$\epsilon \left[(D_x + xy)^2 \partial_x w - 2iy \partial_x w + 2xy^2 w \right] + i(D_y^2 + y) \partial_x w - \lambda \partial_x w = \partial_x f. \quad (5.27)$$

Taking the inner product of (5.27) with $\partial_x w$ yields

$$\begin{aligned} \epsilon & \| (D_x + xy) \partial_x w \|^2 - 2i\epsilon \langle y \partial_x w, \partial_x w \rangle + 2\epsilon \langle xy^2 w, \partial_x w \rangle \\ & + i \| D_y \partial_x w \|^2 + i \langle y \partial_x w, \partial_x w \rangle - \lambda \| \partial_x w \|^2 = \langle \partial_x f, \partial_x w \rangle. \end{aligned} \quad (5.28)$$

Step 7.1. For the case $k = 1/2$, the imaginary part of the above identity reads

$$\| D_y \partial_x w \|^2 + (1 - 2\epsilon) \| y^{1/2} \partial_x w \|^2 + 2\epsilon \operatorname{Im} \langle xy^2 w, \partial_x w \rangle - (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \| \partial_x w \|^2 = \operatorname{Im} \langle \partial_x f, \partial_x w \rangle.$$

Hence, for $0 < \epsilon \leq 1/4$ we have

$$\| y^{1/2} \partial_x w \|^2 \leq 4 \left[(\alpha_0 + \epsilon \operatorname{Im} \mu_1 + 1) \| \partial_x w \|^2 + \epsilon^2 \| x \partial_x w \|^2 + \| y^2 w \|^2 + \| \partial_x f \|^2 \right].$$

We next estimate the first two terms on the right-hand-side of the above inequality using (5.9c) with $k = 0$ and $k = 1$ respectively. The third term is then estimated using (5.9b) with $k = 2$ to obtain

$$\| y^{1/2} \partial_x w \|^2 \leq c \left[\| w \|^2 + \epsilon^{-1} \| (|x|^{3/2} + y^{3/2} + 1) f \|^2 + \| (|x|^{5/2} + y^{3/2} + y^{5/2} + 1) f \|^2 + \| \partial_x f \|^2 \right].$$

Since

$$\| |x|^{3/2} f \|^2 \leq \| (|x|^{5/2} + 1) f \|^2, \quad \| y^{3/2} f \|^2 \leq \| (y^{5/2} + 1) f \|^2,$$

we get

$$\| y^{1/2} \partial_x w \|^2 \leq c \left[\| w \|^2 + \epsilon^{-1} \| (|x|^{5/2} + y^{5/2} + 1) f \|^2 + \| \partial_x f \|^2 \right].$$

Hence (5.9d) is proved for $k = 1/2$.

Step 7.2. For the case $k = 1$, we take the inner product of (5.27) with $y \partial_x w$ to obtain

$$\begin{aligned} \epsilon & \| y^{1/2} (D_x + xy) \partial_x w \|^2 - 2i\epsilon \| y \partial_x w \|^2 + 2\epsilon \langle xy^2 w, y \partial_x w \rangle + i \| y^{1/2} D_y \partial_x w \|^2 \\ & + i \langle \partial_y \partial_x w, \partial_x w \rangle + i \| y \partial_x w \|^2 - \lambda \langle \partial_x w, y \partial_x w \rangle = \langle y^{1/2} \partial_x f, y^{1/2} \partial_x w \rangle. \end{aligned} \quad (5.29)$$

As $w \in D(\mathcal{B}_\epsilon)$ implies $w(x, 0) = 0$, we have that $\partial_x w(x, 0) = 0$. Consequently,

$$\langle \partial_y \partial_x w, \partial_x w \rangle = 0.$$

The imaginary part of (5.29) then reads

$$\begin{aligned} & \| y^{1/2} D_y \partial_x w \|^2 + (1 - 2\epsilon) \| y \partial_x w \|^2 + 2\epsilon \operatorname{Im} \langle xy^2 w, y \partial_x w \rangle \\ & - (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \| y^{1/2} \partial_x w \|^2 = \operatorname{Im} \langle y^{1/2} \partial_x f, y^{1/2} \partial_x w \rangle. \end{aligned}$$

Hence, for $0 < \epsilon \leq 1/4$, we obtain

$$\| y \partial_x w \|^2 \leq 4 \left[(\alpha_0 + \epsilon \operatorname{Im} \mu_1 + 1) \| y^{1/2} \partial_x w \|^2 + \| y^{1/2} \partial_x f \|^2 + \epsilon^2 \| x \partial_x w \|^2 + \| y^3 w \|^2 \right]. \quad (5.30)$$

To estimate $\|y^{1/2}\partial_x w\|^2$ on the right-hand-side we may use (5.9d) for $k = 1/2$. It seems, however, more transparent to use instead the following inequality

$$\begin{aligned} \int_{\mathbb{R}_+^2} y^{n-l} |\partial_x w|^2 dx &\leq \frac{1}{N^{l+1}} \int_{\{y \geq N\}} y^{n+1} |\partial_x w|^2 dx dy + \int_{\{0 \leq y \leq N\}} y^{n-l} |\partial_x w|^2 dx dy \\ &\leq \frac{1}{N^{l+1}} \|y^{(n+1)/2} \partial_x w\|^2 + N^{n-l} \|\partial_x w\|^2, \end{aligned} \quad (5.31)$$

which holds for all $N > 0$. The above inequality, with N being sufficiently large, $n = 1$, and $l = 0$, combined together with (5.30), yields

$$\|y \partial_x w\|^2 \leq c_7 \left[\|\partial_x w\|^2 + \|y^{1/2} \partial_x f\|^2 + \epsilon^2 \|x \partial_x w\|^2 + \|y^3 w\|^2 \right]. \quad (5.32)$$

Combining the above, (5.9b) (for $k = 3$), and (5.9c) (for $k = 0$ and 1) yields the existence of ϵ_0 , c_8 and c_9 , such that, for all $\epsilon \in (0, \epsilon_0]$ we have

$$\begin{aligned} \|y \partial_x w\| &\leq c_8 \left[\|w\| + \epsilon^{-1} \left(\|(|x|^{3/2} + y^{3/2} + 1)f\| + \|y^{1/2} \partial_x f\|^2 \right) \right. \\ &\quad \left. + \epsilon \|w\| + \left(\|(|x|^{5/2} + y^{5/2} + 1)f\| + \|w\| + \|(y^{5/2} + 1)f\| \right) \right] \\ &\leq c_9 \left[\|w\| + \epsilon^{-1} \left(\|(|x|^{5/2} + y^{5/2} + 1)f\| + \|y^{1/2} \partial_x f\| \right) \right], \end{aligned} \quad (5.33)$$

which completes the proof of (5.9d) for $k = 1$.

Step 7.3. Finally, we consider the case $k = (m + 1)/2$. Upon taking the inner product of (5.27) with $y^m \partial_x w$ for an integer $m \geq 1$, we use the identity

$$i \langle D_y^2 \partial_x w, y^m \partial_x w \rangle = i \|D_y(y^{m/2} \partial_x w)\|^2 - \frac{im^2}{4} \|y^{(m/2-1)} \partial_x w\|^2 - m \operatorname{Im} \int_{\mathbb{R}_+^2} y^{m-1} \partial_y \partial_x w \partial_x \bar{w} dx dy,$$

to obtain

$$\begin{aligned} &\epsilon \|y^{m/2} (D_x + xy) \partial_x w\|^2 - 2i\epsilon \|y^{(m+1)/2} \partial_x w\|^2 + 2\epsilon \langle y^{m+2} w, x \partial_x w \rangle + i \|D_y(y^{m/2} \partial_x w)\|^2 \\ &\quad - \frac{im^2}{4} \|y^{m/2-1} \partial_x w\|^2 - m \operatorname{Im} \int_{\mathbb{R}_+^2} y^{m-1} \partial_y \partial_x w \partial_x \bar{w} dx dy + i \|y^{(m+1)/2} \partial_x w\|^2 - \lambda \|y^{m/2} \partial_x w\|^2 \\ &= \langle y^{m/2} \partial_x f, y^{m/2} \partial_x w \rangle. \end{aligned}$$

The imaginary part of the above is given in the form

$$\begin{aligned} &\|D_y(y^{m/2} \partial_x w)\|^2 - \frac{m^2}{4} \|y^{m/2-1} \partial_x w\|^2 + (1 - 2\epsilon) \|y^{(m+1)/2} \partial_x w\|^2 \\ &\quad + 2\epsilon \operatorname{Im} \langle y^{m+2} w, x \partial_x w \rangle - (\alpha_0 + \epsilon \operatorname{Im} \mu_1) \|y^{m/2} \partial_x w\|^2 = \operatorname{Im} \langle y^{m/2} \partial_x f, y^{m/2} \partial_x w \rangle. \end{aligned}$$

Hence, there exists c_{10} , such that, for all $0 < \epsilon \leq 1/4$, we have

$$\begin{aligned} \|y^{(m+1)/2} \partial_x w\|^2 &\leq c_{10} \{ \|y^{m/2-1} \partial_x w\|^2 + \|y^{m+2} w\|^2 + \epsilon^2 \|x \partial_x w\|^2 \\ &\quad + \|y^{m/2} \partial_x w\|^2 + \|y^{m/2} \partial_x f\|^2 \}. \end{aligned} \quad (5.34)$$

To complete the proof we use (5.31) with $n = m$ twice: with $l = 0$ to control $\|y^m \partial_x w\|$, and with $l = 2$ to control $\|y^{m-2} \partial_x w\|$. For sufficiently large N , depending only on c_9 , (5.34) can be converted into the form

$$\|y^{(m+1)/2} \partial_x w\|^2 \leq c_{11} \left[\|\partial_x w\|^2 + \epsilon^2 \|x \partial_x w\|^2 + \|y^{m+2} w\|^2 + \|y^{m/2} \partial_x f\|^2 \right]. \quad (5.35)$$

Using (5.9c) with $k = 0$ and $k = 1$, and (5.9b) with $k = m + 2$, we derive from (5.35) the following estimate

$$\|y^{(m+1)/2}\partial_x w\| \leq c_{12} \left[\|w\| + \epsilon^{-1} \left(\| |x|^{3/2} + y^{3/2} + 1 \right) \|f\| + \left(\| |x|^{5/2} + y^{5/2} + y^{m+3/2} \right) \|f\| + \|y^{m/2}\partial_x f\| \right].$$

Observing that

$$\| |x|^{3/2} f \| \leq \| (|x|^{5/2} + 1) f \|, \quad \| (y^{3/2} + y^{5/2}) f \| \leq 2 \| (y^{m+3/2} + 1) f \|,$$

(we use these inequalities frequently in the sequel without referring to them anymore) we obtain

$$\|y^{(m+1)/2}\partial_x w\| \leq c_{13} \left[\|w\| + \epsilon^{-1} \left(\| |x|^{5/2} + y^{m+3/2} + 1 \right) \|f\| + \|y^{m/2}\partial_x f\| \right]. \quad (5.36)$$

Thus, (5.9d) is true for $k = (m + 1)/2$.

Step 8. Proof of (5.9e). The real part of (5.28) reads

$$\epsilon \| (D_x + xy)\partial_x w \|^2 + 2\epsilon \operatorname{Re} \langle y^2 w, x\partial_x w \rangle - \epsilon \operatorname{Re} \mu_1 \|\partial_x w\|^2 = \operatorname{Re} \langle \partial_x f, \partial_x w \rangle. \quad (5.37)$$

Since

$$\| (D_x + xy)\partial_x w \|^2 = \|\partial_{xx} w\|^2 + \|xy\partial_x w\|^2 + 2\operatorname{Im} \langle xy\partial_x w, \partial_{xx} w \rangle,$$

we obtain

$$\begin{aligned} \|\partial_{xx} w\|^2 &\leq 8\|xy\partial_x w\|^2 + 2\|y^2 w\|^2 + (1 + 2|\mu_1|)\|\partial_x w\|^2 + \epsilon^{-2}\|\partial_x f\|^2 \\ &\leq 2\|y^2 w\|^2 + (1 + 2|\mu_1|)\|\partial_x w\|^2 + 4\|x^2 \partial_x w\|^2 + 4\|y^2 \partial_x w\|^2 + \epsilon^{-2}\|\partial_x f\|^2. \end{aligned}$$

Using the above, (5.9b) with $k = 2$, (5.9c) with $k = 0$ and $k = 2$, and (5.9d) with $k = 2$, we obtain the existence of c and ϵ_0 such that

$$\|\partial_{xx} w\| \leq c \left[\|w\| + \epsilon^{-1} \left(\| |x|^{7/2} + y^{7/2} + 1 \right) \|f\| + \|y^{3/2}\partial_x f\| + \epsilon^{-1} \|\partial_x f\| \right].$$

This verifies (5.9e). ■

It seems worthwhile to note that, if one conducts the various computations in the proof of Lemma 5.2 differently, it is possible to get better estimates for (5.9e). However, all these estimates would fit into the general form

$$\|\partial_{xx}^2 w\| \leq C \left[\|w\| + \epsilon^{-1} p(f) \right],$$

for an appropriately defined norm p on \mathcal{S} .

5.2. Projection on $(\operatorname{span}\{\psi_0\})^\perp$.

We begin by recalling from (2.11) the definition of π_\parallel , the orthogonal projector on the complex 1-dimensional linear subspace generated by ψ_0 . Then, we introduce the complementary projection

$$\pi_\perp = I - \pi_\parallel, \quad (5.38)$$

where I is the identity map. Making use of topological tensor products, we introduce

$$\Pi_\parallel = \pi_\parallel \otimes I \quad \text{and} \quad \Pi_\perp = \pi_\perp \otimes I,$$

which are orthogonal projectors acting on

$$L^2(\mathbb{R}_+, \mathbb{C}) = L^2(\mathbb{R}_+, \mathbb{C}) \widehat{\otimes} L^2(\mathbb{R}, \mathbb{C}).$$

For u in $L^2(\mathbb{R}_+, \mathbb{C})$, we write

$$u_\parallel = \Pi_\parallel u \quad \text{and} \quad u_\perp = \Pi_\perp u.$$

It is easy to show that for all $u \in D(\mathcal{L})$, where \mathcal{L} is the Airy operator defined in (2.10), we have

$$\alpha_0 \|u_{\parallel}\|^2 + \alpha_1 \|u_{\perp}\|^2 \leq \langle \mathcal{L}u, u \rangle, \quad (5.39)$$

where α_0 and α_1 are the lowest and the second eigenvalues of \mathcal{L} , and $\alpha_1 > \alpha_0$.

Given $\epsilon > 0$, $\lambda \in \mathcal{C}_{\epsilon, \rho}$ and $f \in \mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$, let $w \in D(\mathcal{B}_{\epsilon})$ be the solution of (3.13). Write

$$w = w_{\parallel} + w_{\perp}. \quad (5.40)$$

We next show that for a suitable choice of ρ , w_{\perp} is small, compared with w_{\parallel} , if ϵ and f are sufficiently small.

Lemma 5.3. *There exist positive constants C , k_0 and ϵ_0 such that, for any $\epsilon \in (0, \epsilon_0]$, if $\lambda = \lambda(\mu_1)$ is given by (5.8), and if $(w, f) \in D(\mathcal{B}_{\epsilon}) \times \mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$ satisfies (3.13), then we have:*

$$\|w_{\perp}\|^2 \leq \frac{1}{\alpha_1 - \alpha_0 - 2\epsilon|\mu_1|} \left[\epsilon|\mu_1| \|w_{\parallel}\|^2 + \|w\| \|f\| \right], \quad (5.41a)$$

$$\|x^2 w_{\perp}\|^2 \leq C \left[\epsilon \|w\|^2 + \epsilon^{-1} \|(1 + x^2)f\|^2 \right], \quad (5.41b)$$

$$\|x \partial_x w_{\perp}\|^2 \leq C \left[\epsilon \|w\|^2 + \epsilon^{-1} p_{k_0}(f)^2 \right], \quad (5.41c)$$

where p_{k_0} is a norm in $\mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$ defined in (3.6).

Proof. Step 1. We prove (5.41a). Let w be the solution of (3.13). Recalling (5.8) we take the inner product of (3.13) with w and then obtain by considering the imaginary part of the new expression:

$$\langle (\mathcal{L} - \mu_0)w, w \rangle = \text{Im} \langle f, w \rangle, \quad (5.42)$$

with

$$\mu_0 = \text{Im} \lambda = \alpha_0 + \epsilon \text{Im} \mu_1 \quad (5.43)$$

It follows from this and (5.39) that,

$$(\alpha_1 - \alpha_0 - \epsilon \text{Im} \mu_1) \|w_{\perp}\|^2 - \epsilon \text{Im} \mu_1 \|w_{\parallel}\|^2 \leq \|f\| \|w\|,$$

from which we deduce (5.41a) for small ϵ .

Step 2. We prove (5.41b). Taking the inner product of (3.13) with $x^{2k}w$, where k is any positive integer, and computing as in (5.3), we obtain

$$\begin{aligned} \langle f, x^{2k}w \rangle &= \langle \mathcal{B}_{\epsilon}w - \lambda w, x^{2k}w \rangle = \langle \epsilon(D_x + xy)^2 w, w \rangle + \langle (i\mathcal{L} - \lambda)w, x^{2k}w \rangle \\ &= \epsilon \|x^k(D_x + xy)w\|^2 + 2ki\epsilon \langle (D_x + xy)w, x^{2k-1}w \rangle + \langle (i\mathcal{L} - \lambda)w, x^{2k}w \rangle \\ &= \epsilon \|x^k(D_x + xy)w\|^2 + 2ki\epsilon \langle (D_x + xy)(x^k w), x^{k-1}w \rangle - 2k^2\epsilon \|x^{k-1}w\|^2 \\ &\quad + \langle (i\mathcal{L} - \lambda)(x^k w), x^k w \rangle. \end{aligned}$$

Combining the imaginary part of the above identity and (5.42) yields

$$2k\epsilon \text{Re} \langle (D_x + xy)(x^k w), x^{k-1}w \rangle + \langle (\mathcal{L} - \mu_0)(x^k w), x^k w \rangle = \text{Im} \langle x^k f, x^k w \rangle.$$

We next make the obvious observation that

$$\pi_{\parallel}(x^k w) = x^k \pi_{\parallel} w = x^k w_{\parallel}, \quad \pi_{\perp}(x^k w) = x^k \pi_{\perp} w = x^k w_{\perp}.$$

Thus, from (5.39) it follows that,

$$\begin{aligned}
& \alpha_0 \|x^k w_{\parallel}\|^2 + \alpha_1 \|x^k w_{\perp}\|^2 \leq \langle \mathcal{L}(x^k w), x^k w \rangle \\
& \leq \mu_0 \|x^k w\|^2 + \|x^k f\| \|x^k w\| + 2k\epsilon \|(D_x + xy)(x^k w)\| \|x^{k-1} w\| \\
& \leq \mu_0 \|x^k w_{\parallel}\|^2 + \mu_0 \|x^k w_{\perp}\|^2 + \frac{1}{2}\epsilon^{-1} \|x^k f\|^2 + \frac{\epsilon}{2} \|x^k w\|^2 \\
& \quad + k\epsilon \|x^{k-1} w\|^2 + k\epsilon \|(D_x + xy)(x^k w)\|^2.
\end{aligned} \tag{5.44}$$

To control the last term in the right side of (5.44) we use (5.13) with $m = 2k$ to obtain the existence of c_1 for which we have

$$\begin{aligned}
& k\epsilon \int_{\mathbb{R}_+^2} |(D_x + xy)(x^k w)|^2 dx dy \\
& \leq k \operatorname{Re} \lambda \|x^k w\|^2 + k\epsilon \|(x^k + kx^{k-1})w\|^2 + k \|x^k w\| \|x^k f\| \\
& = k\epsilon(\lambda_1 + \mu_1) \|x^k w\|^2 + 2k\epsilon \|x^k w\|^2 + 2k^2\epsilon \|x^{k-1} w\|^2 + \frac{k}{2}\epsilon \|x^k w\|^2 + \frac{k}{2}\epsilon^{-1} \|x^k f\|^2 \\
& \leq c_1\epsilon (\|x^k w\|^2 + \|x^{k-1} w\|^2) + \frac{k}{2}\epsilon^{-1} \|x^k f\|^2.
\end{aligned}$$

Substituting it back into (5.44) we derive the existence of c_2 such that

$$(\alpha_0 - \mu_0) \|x^k w_{\parallel}\|^2 + (\alpha_1 - \mu_0) \|x^k w_{\perp}\|^2 \leq c_2\epsilon (\|x^k w\|^2 + \|x^{k-1} w\|^2) + c_2\epsilon^{-1} \|x^k f\|^2.$$

By (5.43) and (5.9a) we then have

$$\begin{aligned}
& (\alpha_1 - \alpha_0) \|x^k w_{\perp}\|^2 \\
& \leq \epsilon \operatorname{Im} \mu_1 (\|x^k w_{\parallel}\|^2 + \|x^k w_{\perp}\|^2) + c_2\epsilon (\|x^k w\|^2 + \|x^{k-1} w\|^2) + c_2\epsilon^{-1} \|x^k f\|^2 \\
& \leq c_3\epsilon (\|x^k w\|^2 + \|x^{k-1} w\|^2) + c_3\epsilon^{-1} \|x^k f\|^2 \\
& \leq c_4\epsilon \left[\|w\|^2 + \epsilon^{-2} (|x|^{k-1/2} + |x|^{k-3/2} + 1) f^2 \right] + c_2\epsilon^{-1} \|x^k f\|^2 \\
& = c_4\epsilon \|w\|^2 + (c_2 + c_4)\epsilon^{-1} \|x^k + |x|^{k-1/2} + |x|^{k-3/2} + 1\| f\|^2.
\end{aligned}$$

Hence

$$(\alpha_1 - \alpha_0) \|x^k w_{\perp}\|^2 \leq c_5\epsilon \|w\|^2 + c_5\epsilon^{-1} \|(x^k + 1)f\|^2, \tag{5.45}$$

which with $k = 2$ yields (5.41b).

Step 3. Proof of (5.41c). We first differentiate (3.13) with respect to x (see also (5.27)) to obtain

$$(i\mathcal{L} - \lambda)(\partial_x w) + \epsilon \partial_x (D_x + xy)^2 w = \partial_x f.$$

Then, taking the inner product of the above identity with $x^2 \partial_x w$ yields

$$\langle (i\mathcal{L} - \lambda)(x \partial_x w), x \partial_x w \rangle + \epsilon \langle \partial_x (D_x + xy)^2 w, x^2 \partial_x w \rangle = \langle \partial_x f, x^2 \partial_x w \rangle.$$

The imaginary part, with the aid of (5.43), receives the form

$$\begin{aligned}
& \langle (\mathcal{L} - \mu_0)(x \partial_x w), x \partial_x w \rangle = \operatorname{Im} \langle \partial_x f, x^2 \partial_x w \rangle - \epsilon \operatorname{Im} \langle \partial_x (D_x + xy)^2 w, x^2 \partial_x w \rangle \\
& = \operatorname{Im} \langle \partial_x f, x^2 \partial_x w \rangle + 2\epsilon \|xy^{1/2} \partial_x w\|^2 - \epsilon \operatorname{Im} \langle y^2 w, x^3 \partial_x w \rangle + \epsilon \operatorname{Im} \langle (D_x + xy)^2 \partial_x w, x^2 \partial_x w \rangle.
\end{aligned}$$

Note that

$$\langle (D_x + xy)^2 \partial_x w, x^2 \partial_x w \rangle = \|x(D_x + xy) \partial_x w\|^2 + 2\langle \partial_{xx} w, x \partial_x w \rangle + 2i \|xy^{1/2} \partial_x w\|^2.$$

Consequently, we have

$$\begin{aligned} \langle (\mathcal{L} - \mu_0)(x\partial_x w), x\partial_x w \rangle &= \text{Im} \langle \partial_x f, x^2 \partial_x w \rangle - \epsilon \text{Im} \langle y^2 w, x^3 \partial_x w \rangle \\ &\quad + 2\epsilon \text{Im} \langle \partial_{xx} w, x\partial_x w \rangle + 2\epsilon \|xy^{1/2} \partial_x w\|^2. \end{aligned}$$

Combining the above with (5.39) we obtain

$$\begin{aligned} &(\alpha_0 - \mu_0) \|x\partial_x w_{\parallel}\|^2 + (\alpha_1 - \mu_0) \|x\partial_x w_{\perp}\|^2 \\ &\leq \langle (\mathcal{L} - \mu_0)(x\partial_x w), x\partial_x w \rangle \\ &\leq \|\partial_x f\| \|x^2 \partial_x w\| + \epsilon \|y^2 w\| \|x^3 \partial_x w\| + 2\epsilon |\langle \partial_{xx} w, x\partial_x w \rangle| + 2\epsilon \|xy^{1/2} \partial_x w\|^2, \end{aligned}$$

which together with (5.43) yields

$$\begin{aligned} &(\alpha_1 - \mu_0) \|x\partial_x w_{\perp}\|^2 \\ &\leq C\epsilon [\|x^2 \partial_x w_{\parallel}\|^2 + \|x\partial_x w\|^2 + \|y^2 w\|^2 + \|x^3 \partial_x w\|^2 + |\langle \partial_{xx} w, x\partial_x w \rangle| + \|xy^{1/2} \partial_x w\|^2] \\ &\quad + \epsilon^{-1} \|\partial_x f\|^2 \\ &\leq C\epsilon [\|y^2 w\|^2 + \|x\partial_x w\|^2 + \|x^2 \partial_x w\|^2 + \|x^3 \partial_x w\|^2 + \|y\partial_x w\|^2 + \|\partial_{xx} w\|^2] + \epsilon^{-1} \|x\partial_x f\|^2. \end{aligned}$$

Using (5.9b) with $k = 1$, (5.9c) with $k = 1, 2, 3$, (5.9d) with $k = 1$, and (5.9e) we obtain, for all $0 < \epsilon \leq 1/4$, that

$$\begin{aligned} &(\alpha_1 - \mu_0) \|x\partial_x w_{\perp}\|^2 \\ &\leq C\epsilon \{ \|w\|^2 + \epsilon^{-2} \|(1 + |x|^{9/2} + y^{9/2})f\|^2 + \|(y^{1/2} + y^{3/2})\partial_x f\|^2 + \epsilon^{-2} \|\partial_x f\|^2 \} + \epsilon^{-1} \|x\partial_x f\|^2. \end{aligned}$$

Hence, (5.41c) is verified for $0 < \epsilon \leq 1/4$, with

$$p(f)^2 = \|(1 + |x|^{9/2} + y^{9/2})f\|^2 + \|(1 + |x| + y^{3/2})\partial_x f\|^2.$$

■

We are now ready for the final stage of Step 4 of the proof for the main results stated in Section 3, in which we obtain an estimate of the norm of the solution $w = (\mathcal{B}_{\epsilon} - \lambda)^{-1} f$ of (3.13) for $\lambda \in \mathcal{C}_{\epsilon, \rho}$ and $f \in \mathcal{S}$. Recall that every λ on the circle $\mathcal{C}_{\epsilon, \rho}$ obeys the form

$$\lambda = \lambda(\delta) = \gamma_{\epsilon} + \epsilon\delta(\rho, \phi, \epsilon), \quad (5.46)$$

where γ_{ϵ} is given in (3.4), and

$$\delta(\rho, \phi, \epsilon) := \rho \exp(i\phi) \epsilon^{1/2}, \quad \phi \in [0, 2\pi). \quad (5.47)$$

Lemma 5.4. *There exist $k_0 \in \mathbb{N}$, $\rho_0 > 0$, and $C > 0$, such that for any w satisfying (3.13) with $\lambda \in \mathcal{C}_{\epsilon, \rho}$ and $\rho > \rho_0$ it holds that*

$$\exists \epsilon_0(\rho) : \quad 0 < \epsilon < \epsilon_0(\rho) \implies \|w\| \leq C(\epsilon|\delta|)^{-1} p_{k_0}(f), \quad (5.48)$$

where p_{k_0} is the norm in $\mathcal{S}(\overline{\mathbb{R}_+^2}, \mathbb{C})$ given in Lemma 5.3.

Proof. Step 1. Let

$$w_1(x) = \int_0^{+\infty} w(x, y) \psi_0(y) dy.$$

Then,

$$w_{\parallel}(x, y) = w_1(x) \psi_0(y), \quad \|w_{\parallel}\| = \|w_1\|_{L^2(\mathbb{R})}.$$

Taking the inner product of (3.13) with ψ_0 in $L^2(\mathbb{R}_+, \mathbb{C})$ with respect to the variable y , and dividing the resulting equation by ϵ we obtain

$$D_x^2 w_1 - 2ix\partial_x \langle w, y\psi_0 \rangle_y - i \langle w, y\psi_0 \rangle_y + x^2 \langle w, y^2\psi_0 \rangle_y - \frac{\lambda - i\alpha_0}{\epsilon} w_1 = \frac{1}{\epsilon} \langle f, w \rangle_y, \quad (5.49)$$

where $\langle \cdot, \cdot \rangle_y$ denotes the inner product in $L^2(\mathbb{R}_+, \mathbb{C})$. Having in mind the definition (4.9)-(4.10) of the operator \mathcal{K}_0 , we can rewrite $D_x^2 \phi$, for any smooth ϕ , in the form

$$D_x^2 \phi = \mathcal{K}_0 \phi + 2i\alpha x \partial_x \phi + i\alpha \phi - \beta^2 x^2 \phi. \quad (5.50)$$

Therefore,

$$\begin{aligned} & D_x^2 w_1 - 2ix\partial_x \langle w, y\psi_0 \rangle_y - i \langle w, y\psi_0 \rangle_y + x^2 \langle w, y^2\psi_0 \rangle_y \\ &= \mathcal{K}_0 w_1 - 2ix\partial_x [\langle w, y\psi_0 \rangle_y - \alpha w_1] - i [\langle w, y\psi_0 \rangle_y - \alpha w_1] + x^2 [\langle w, y^2\psi_0 \rangle_y - \beta^2 w_1] \\ &= \mathcal{K}_0 w_1 - 2ix\partial_x \langle w_\perp, y\psi_0 \rangle_y - i \langle w_\perp, y\psi_0 \rangle_y + x^2 \langle w_\perp, y^2\psi_0 \rangle_y, \end{aligned} \quad (5.51)$$

where we have used the identities:

$$\begin{aligned} \langle w, y\psi_0 \rangle_y - \alpha w_1 &= \int_0^{+\infty} y\psi_0(w - w_1(x)\psi_0(y))dy = \langle w - w_\parallel, y\psi_0 \rangle_y = \langle w_\perp, y\psi_0 \rangle_y, \\ \langle w, y^2\psi_0 \rangle_y - \beta^2 w_1 &= \int_0^{+\infty} y^2\psi_0(w - w_1\psi_0)dy = \langle w - w_\parallel, y^2\psi_0 \rangle_y = \langle w_\perp, y^2\psi_0 \rangle_y. \end{aligned}$$

We then combine (5.49) and (5.51), using (5.46) and (5.47) to get

$$(\mathcal{K}_0 - \lambda_1 - \delta)w_1 = g, \quad (5.52)$$

where

$$g = 2ix \langle \partial_x w_\perp, y\psi_0 \rangle_y + i \langle w_\perp, y\psi_0 \rangle_y - x^2 \langle w_\perp, y^2\psi_0 \rangle_y + \epsilon^{-1} \langle f, \psi_0 \rangle_y.$$

Step 2. We next estimate $\|g\|$. Since

$$\begin{aligned} \|x \langle \partial_x w_\perp, y\psi_0 \rangle_y\|_{L^2(\mathbb{R})}^2 &\leq \|x \|y\psi_0\|_{L^2(\mathbb{R}_+)} \|\partial_x w_\perp\|_{L^2(\mathbb{R}_+)}\|_{L^2(\mathbb{R})}^2 \\ &\leq \|y\psi_0\|_{L^2(\mathbb{R}_+)}^2 \|x \partial_x w_\perp\|^2 \leq C \|x \partial_x w_\perp\|^2, \end{aligned}$$

and since there exists $C > 0$ such that

$$\|x^2 \langle y^2\psi_0, w_\perp \rangle_y\|_{L^2(\mathbb{R})}^2 \leq C \|x^2 w_\perp\|^2,$$

we have

$$\|g\|_{L^2(\mathbb{R})}^2 \leq C \left[\|x \partial_x w_\perp\|^2 + \|w_\perp\|^2 + \|x^2 w_\perp\|^2 + \epsilon^{-2} \|f\|^2 \right]. \quad (5.53)$$

Consequently, by (5.41) we have, for some k_0 ,

$$\|g\|_{L^2(\mathbb{R})}^2 \leq C \left[\epsilon \|w\|^2 + \epsilon^{-2} p_{k_0}(f)^2 \right]. \quad (5.54)$$

Step 3. As the operator \mathcal{K}_0 is self-adjoint we obtain from (5.52) and the spectral theorem [16] that for any fixed $\rho > 0$ there exists $\epsilon_0(\rho) > 0$ such that for all $0 < \epsilon < \epsilon_0(\rho)$ we have

$$\|w_1\|_{L^2(\mathbb{R})} \leq 2 \frac{\|g\|_{L^2(\mathbb{R})}}{|\delta|}. \quad (5.55)$$

Substituting (5.54) into (5.55) we obtain for some $k_0 \in \mathbb{N}$,

$$\|w_1\|_{L^2(\mathbb{R})} \leq C_1 |\delta|^{-1} \left[\epsilon^{1/2} \|w\| + \epsilon^{-1} p_{k_0}(f) \right]. \quad (5.56)$$

By (5.41a), there exists $C_2 > 0$ such that, for small ϵ ,

$$\|w\|^2 \leq C_2 \left[\|w_{\parallel}\|^2 + \|w\| \|f\| \right],$$

and hence, noting that $\|w_1\|_{L^2(\mathbb{R})} = \|w_{\parallel}\|$, we have that

$$\|w_1\|_{L^2(\mathbb{R})}^2 \geq \frac{1}{2C_2} \left[\|w\|^2 - C_2^2 \|f\|^2 \right]. \quad (5.57)$$

Using the above inequality together with (5.56), we obtain the existence of k_0 and C_3 such that

$$\|w\|^2 \leq C_2^2 \|f\|^2 + 2C_2 \|w_1\|_{L^2(\mathbb{R})}^2 \leq C_3 \delta^{-2} \left[\epsilon \|w\|^2 + \epsilon^{-2} p_{k_0}(f)^2 \right], \quad (5.58)$$

which is valid for all $\epsilon \in (0, \epsilon_0(\rho))$.

Let then

$$\rho_0 = \sqrt{2C_3}.$$

For a given $\rho > \rho_0$, we obtain from (5.58) that for all $\epsilon \in (0, \epsilon_0(\rho))$, there exist $k_0 \in \mathbb{N}$ and $C_4 > 0$ such that

$$\|w\| \leq C_4 (\epsilon \delta)^{-1} p_{k_0}(f).$$

■

As an immediate consequence of Lemma 5.4 we obtain

Proposition 5.5. *Let w satisfy (3.13) with $f \equiv 0$ and $\lambda \in \mathcal{C}_{\epsilon, \rho}$, where ϵ is small enough. Then $w \equiv 0$.*

Proof of Proposition 3.4.

(3.11) follows from (5.48) with $|\delta| = \rho \epsilon^{1/2}$. ■

The other consequences of Lemma 5.4 have been described at the end of Section 3.

6. EXISTENCE, UPPER AND LOWER BOUNDS

It is now possible to state the following existence result:

Proposition 6.1. *Let*

$$\gamma_{\epsilon} = i\alpha_0 + \epsilon\lambda_1, \quad (6.1)$$

where α_0 is the first eigenvalue of the Airy operator \mathcal{L} and λ_1 is the first eigenvalue of the operator \mathcal{K}_0 defined in (4.9). Then there exist $\epsilon_0 > 0$ and $C > 0$ such that, for all $0 < \epsilon < \epsilon_0$,

$$\sigma(\mathcal{B}_{\epsilon}) \cap B(\gamma_{\epsilon}, C\epsilon^{3/2}) \neq \emptyset. \quad (6.2)$$

Proof. The proof follows immediately from (3.10) and (5.48). ■

By (5.48) it follows that $\mathcal{B}_{\epsilon} - \lambda$ is injective for $\lambda \in \mathcal{C}(\epsilon, \rho)$ with ϵ small enough. As \mathcal{B}_{ϵ} has a compact resolvent it immediately follows that $\lambda \notin \sigma(\mathcal{B}_{\epsilon})$.

Coming back to our initial operator \mathcal{A}_c^+ we deduce from Proposition 6.1 the following corollary:

Corollary 6.2. *The operator \mathcal{A}_c^+ defined in the introduction admits at least one eigenvalue $\mu = \mu(c)$ having the following expansion as $|c| \rightarrow +\infty$:*

$$\mu(c) = c^{2/3} \exp(i\frac{\pi}{3})\alpha_0 + \lambda_1 \exp(-i\frac{\pi}{6})c^{-1/3} + \mathcal{O}(c^{-5/6}). \quad (6.3)$$

Clearly, (6.3) establishes the upper bound in (1.18). To obtain the lower bound we need to prove first Proposition 3.5.

Proof of Proposition 3.5. The resolvent of \mathcal{B}_ϵ being compact, it is sufficient to show that there are no eigenvalues in $D_{b,\epsilon}$. Let then $\lambda \in D_{b,\epsilon}$, where $D_{b,\epsilon}$ is given by (3.14), and let $w \in D(\mathcal{B}_\epsilon)$ satisfy

$$(\mathcal{B}_\epsilon - \lambda)w = 0.$$

We note first that, for fixed b and sufficiently small $\epsilon > 0$, both (5.9) and (5.41) hold for $\lambda \in D_{b,\epsilon}$. Furthermore, despite the fact that Lemma 5.4 is stated for $\lambda \in \mathcal{C}_{\epsilon,\rho}$ (see (5.46), (5.47)), we can still use (5.54), (5.57), and even (5.52) as long as we appropriately modify the definition of δ . More precisely, $\lambda = \gamma_\epsilon + \epsilon\delta \in D_{b,\epsilon}$ means that instead of (5.47) we have

$$-\lambda_1 < \operatorname{Re} \delta \leq -b\epsilon^{1/2}, \quad 0 < \operatorname{Im} \delta \leq 3\lambda_1.$$

We can thus take the inner product with respect to x , in $L^2(\mathbb{R}, \mathbb{C})$, of (5.52) with w_1 , defined in the proof of Lemma 5.4, to obtain

$$-\operatorname{Re} \delta \|w_1\|_{L^2(\mathbb{R})}^2 \leq \operatorname{Re} \langle (\mathcal{K}_0 - \lambda_1)w_1, w_1 \rangle_x - \operatorname{Re} \delta \|w_1\|_{L^2(\mathbb{R})}^2 = \langle g, w_1 \rangle_x \leq \|g\|_{L^2(\mathbb{R})} \|w_1\|_{L^2(\mathbb{R})},$$

where g was defined after (5.52). By (5.54), with $f = 0$, it then follows that

$$b\epsilon^{1/2} \|w_1\|_{L^2(\mathbb{R})} \leq C\epsilon^{1/2} \|w\|.$$

Using (5.57) with $f = 0$ then yields

$$\frac{b}{\sqrt{2C_2}} \|w\| \leq C \|w\|.$$

For sufficiently large b we must then have $w \equiv 0$, which means that λ is not an eigenvalue. ■

7. DECAY OF THE SEMI-GROUP

In this section we apply a technique which was applied first in [13] to obtain stability of the normal states for the linearized one-dimensional Ginzburg-Landau system, in the absence of any magnetic field (induced or applied). The linear operator in this case is the Dirichlet realization of the complex Airy operator in \mathbb{R}_+ [14], which we denote here by \mathcal{Q} . In [3], it has been shown that when λ lies to the left of $\sigma(\mathcal{Q})$, then

$$\|e^{-t(\mathcal{Q}-\lambda)}u_0\|_{L^2(\mathbb{R}_+)} \xrightarrow[t \rightarrow +\infty]{} 0,$$

when u_0 belongs to a certain dense set in $L^2(\mathbb{R}_+)$. In [13] the validity of the above has been extended to any $u_0 \in L^2(\mathbb{R}_+)$. We use the technique in [13] to obtain a similar result, but this time for the operator \mathcal{A}_c^+ , i.e., we prove Theorem 1.3. This desired outcome is a consequence of the following proposition.

Recall the definition of $\mu_m(c)$ in (1.17).

Proposition 7.1. *For any ω satisfying*

$$\omega < \mu_m, \tag{7.1}$$

there exists $M_\omega > 0$ such that,

$$\|G(t)\|_{\mathcal{L}(L^2(\mathbb{R}_+^2, \mathbb{C}))} \leq M_\omega \exp(-\omega t). \tag{7.2}$$

Furthermore, for any $\omega > \mu_m$ we have that

$$\|e^{\omega t}G(t)\|_{\mathcal{L}(L^2(\mathbb{R}_+^2, \mathbb{C}))} \geq \exp\{(\omega - \mu_m)t\} \tag{7.3}$$

Note that by (2.9) we have that

$$\inf_{z \in \sigma(\mathcal{A}_c^+)} \operatorname{Re} z \geq \frac{1}{2}|c|^{2/3}\alpha_0.$$

Furthermore, for a sufficiently large value of c , in view of (1.18), we can approximate μ_m using (6.3). In the unlikely event that for some value of c , the spectrum of \mathcal{A}_c^+ is empty we may set $\mu_m = +\infty$, i.e., (7.2) would hold for all $\omega \in \mathbb{R}$, and (7.3) would become redundant.

We prove (7.2) with \mathcal{A}_c^+ replaced by \mathcal{P} defined in (2.1) since they are unitarily equivalent by (2.1). In order to make use of the Gearhart-Prüss Theorem (see the Appendix) we need the following lemma:

Lemma 7.2. *Let μ_m be the number defined in (7.1). For any real number α satisfying $\alpha < \mu_m$, there exists a constant C , such that for all λ with $\operatorname{Re} \lambda \leq \alpha$ we have*

$$\|(\mathcal{P} - \lambda)^{-1}\| \leq C. \quad (7.4)$$

Proof. Without loss of generality we assume $\alpha > 0$. Let then λ be such that $\operatorname{Re} \lambda < \alpha$. Clearly, λ lies outside the spectrum of \mathcal{A}_c^+ , and hence outside the spectrum of \mathcal{P} as well. Consequently, once we manage to prove boundedness of the resolvent $(\mathcal{P} - \lambda)^{-1}$ as $|\lambda| \rightarrow +\infty$ (for $\operatorname{Re} \lambda < \alpha$), then the lemma will be proved.

We distinguish between three different subdomains of \mathbb{C} :

- (1) $\operatorname{Re} \lambda \leq -\alpha$.
- (2) $\operatorname{Im} \lambda \rightarrow -\infty$, and $\operatorname{Re} \lambda \in (-\alpha, \alpha)$.
- (3) $\operatorname{Im} \lambda \rightarrow +\infty$, and $\operatorname{Re} \lambda \in (-\alpha, \alpha)$.

In the first region we use the inequality

$$\operatorname{Re} \langle (\mathcal{P} - \lambda)u, u \rangle = \|(D_x + yx)u\|^2 + \|D_y u\|^2 - \operatorname{Re} \lambda \|u\|^2 \geq (E_0^* - \operatorname{Re} \lambda) \|u\|^2, \quad (7.5)$$

where E_0^* was given in (1.14), to obtain that

$$\|(\mathcal{P} - \lambda)^{-1}\| \leq \frac{1}{E_0^* + \alpha}.$$

In the second case we use the inequality

$$\operatorname{Im} \langle (\mathcal{P} - \lambda)u, u \rangle = \int_{\mathbb{R}_+^2} (y - \operatorname{Im} \lambda) |u|^2 \geq -\operatorname{Im} \lambda \|u\|^2, \quad (7.6)$$

yielding immediately that

$$\|(\mathcal{P} - \lambda)^{-1}\| \leq \frac{1}{|\operatorname{Im} \lambda|}.$$

Consequently, it remains necessary to control the norm of the resolvent only in the third case.

Next, we prove the lemma in the third case. To this end, we borrow ideas from semi-classical analysis. The main idea of the proof is to approximate $(\mathcal{P} - \lambda)^{-1}$ by a sum of two different operators: one of them should serve as a good approximation when applied to functions supported near the boundary, while the other one should take care of functions whose support lies far away from the boundary.

The first component in the above decomposition is thus derived from the Dirichlet realization of the differential operator

$$(D_x + yx)^2 + D_y^2 + i(y - \operatorname{Im} \lambda) - \operatorname{Re} \lambda,$$

for (x, y) in the band

$$\mathcal{S} = \mathbb{R} \times \left(0, \frac{\operatorname{Im} \lambda}{2}\right).$$

Denote the resolvent of this operator by

$$\mathcal{R}_1(\lambda) : L^2(\mathcal{S}, \mathbb{C}) \rightarrow H_{\text{mag}}^1(\mathcal{S}, \mathbb{C}).$$

Since (7.6) still holds for the above Dirichlet realization, we easily obtain that

$$\|\mathcal{R}_1(\lambda)\| \leq \frac{2}{\text{Im } \lambda}. \quad (7.7)$$

Furthermore, by (7.5), which remains valid for the problem in \mathcal{S} , we have, for $u = \mathcal{R}_1 f$,

$$\begin{aligned} \|D_y \mathcal{R}_1(\lambda) f\|^2 &= \|D_y u\|^2 \leq \|(\mathcal{P} - \lambda)u\| \|u\| + \text{Re } \lambda \|u\|^2 \\ &\leq \|f\| \|\mathcal{R}_1(\lambda) f\| + |\alpha| \|\mathcal{R}_1(\lambda) f\|^2 \leq \left(\frac{2}{\text{Im } |\lambda|} + \frac{4|\alpha|}{|\text{Im } \lambda|^2} \right) \|f\|^2. \end{aligned}$$

Hence

$$\|D_y \mathcal{R}_1(\lambda)\| \leq \left[\frac{2}{|\text{Im } \lambda|} \right]^{1/2} + \frac{2\alpha^{1/2}}{|\text{Im } \lambda|}. \quad (7.8)$$

Far from the boundary, we attempt to approximate the resolvent of the same differential operator on \mathbb{R}^2 , neglecting the boundary effect. Denote this resolvent by $\mathcal{R}_2(\lambda)$. Recall from [4] that the norm $\|\mathcal{R}_2(\lambda)\|$ is independent of $\text{Im } \lambda$ because the problem in \mathbb{R}^2 is invariant to translations in y variable. Since $\mathcal{R}_2(\lambda)$ is an entire function in λ , we easily obtain a uniform bound on $\|\mathcal{R}_2(\lambda)\|$ for $\text{Re } \lambda \in [-\alpha, +\alpha]$. Hence,

$$\|\mathcal{R}_2(\lambda)\| \leq C_1(\alpha). \quad (7.9)$$

We now use a partition of unity in the y variable in order to construct an approximate inverse $\mathcal{R}^{\text{app}}(\lambda)$ for $\mathcal{P} - \lambda$. We shall then prove that the difference between the approximation and the exact resolvent is well controlled as $\text{Im } \lambda \rightarrow +\infty$. Define then the following pair of cutoff functions in $C^\infty(\mathbb{R}_+, [0, 1])$:

$$\begin{aligned} \phi(t) &= 1 \text{ on } [0, 3/8], \quad \phi = 0 \text{ on } [1/2, +\infty), \\ \psi(t) &= 1 \text{ on } [0, 1/4], \quad \psi = 0 \text{ on } [3/8, +\infty), \end{aligned}$$

and then set

$$\phi_\lambda(y) = \phi\left(\frac{y}{\text{Im } \lambda}\right), \quad \psi_\lambda(y) = \psi\left(\frac{y}{\text{Im } \lambda}\right).$$

The approximate inverse $\mathcal{R}^{\text{app}}(\lambda)$ is then constructed as

$$\mathcal{R}^{\text{app}}(\lambda) = \phi_\lambda \mathcal{R}_1(\lambda) \psi_\lambda + \mathcal{R}_2(\lambda)(1 - \psi_\lambda), \quad (7.10)$$

where ϕ_λ and ψ_λ denote the operators of multiplication by the functions $\phi_\lambda(y)$ and $\psi_\lambda(y)$. Thus,

$$\begin{aligned} \phi_\lambda &: D(\mathcal{R}_1^{-1}) \rightarrow D(\mathcal{P}), \\ \psi_\lambda &: L^2(\mathbb{R}_+^2, \mathbb{C}) \rightarrow L^2(\mathcal{S}, \mathbb{C}), \\ 1 - \psi_\lambda &: L^2(\mathbb{R}_+^2, \mathbb{C}) \rightarrow L^2(\mathbb{R}^2, \mathbb{C}). \end{aligned}$$

From (7.7) and (7.9) we get, for sufficiently large $\text{Im } \lambda$,

$$\|\mathcal{R}^{\text{app}}(\lambda)\| \leq C_3(\alpha). \quad (7.11)$$

Note that

$$\phi_\lambda \psi_\lambda = \psi_\lambda, \quad (7.12)$$

and that

$$|\phi'_\lambda(y)| \leq \frac{C}{|\text{Im } \lambda|}, \quad |\phi''_\lambda(y)| \leq \frac{C}{|\text{Im } \lambda|^2}. \quad (7.13)$$

Next, we apply $\mathcal{P} - \lambda$ to \mathcal{R}^{app} to obtain that :

$$(\mathcal{P} - \lambda)\mathcal{R}^{\text{app}}(\lambda) = I + [\mathcal{P}, \phi_\lambda]\mathcal{R}_1(\lambda) \quad (7.14)$$

where I is the identity operator, and

$$[\mathcal{P}, \phi_\lambda] \equiv \mathcal{P}\phi_\lambda - \phi_\lambda\mathcal{P} = [D_y^2, \phi_\lambda] = -\frac{2i}{\text{Im } \lambda}\phi'_y\left(\frac{y}{\text{Im } \lambda}\right)D_y - \frac{1}{(\text{Im } \lambda)^2}\phi''_y\left(\frac{y}{\text{Im } \lambda}\right). \quad (7.15)$$

Here we have used (7.10), (7.12) and the fact that

$$(\mathcal{P} - \lambda)\mathcal{R}_1(\lambda)\psi_\lambda u = \psi_\lambda u, \quad \forall u \in L^2(\mathbb{R}_+^2, \mathbb{C}).$$

Using (7.7), (7.8), and (7.15) we then easily obtain, for $\text{Im } \lambda$ large enough,

$$\|[\mathcal{P}, \phi_\lambda]\mathcal{R}_1(\lambda)\| \leq \frac{C_4(\alpha)}{|\text{Im } \lambda|^{3/2}}. \quad (7.16)$$

Hence, if $|\text{Im } \lambda|$ is sufficiently large then $I + [\mathcal{P}, \phi_\lambda]\mathcal{R}_1(\lambda)$ is invertible in $\mathcal{L}(L^2(\mathbb{R}_+^2, \mathbb{C}))$, and

$$\|(I + [\mathcal{P}, \phi_\lambda]\mathcal{R}_1(\lambda))^{-1}\| \leq C_5(\alpha). \quad (7.17)$$

Finally, since

$$(\mathcal{P} - \lambda)^{-1} = \mathcal{R}^{\text{app}}(\lambda) \circ (I + [\mathcal{P}, \phi_\lambda]\mathcal{R}_1(\lambda))^{-1},$$

we have that

$$\|(\mathcal{P} - \lambda)^{-1}\| \leq \|\mathcal{R}^{\text{app}}(\lambda)\| \|(I + [\mathcal{P}, \phi_\lambda]\mathcal{R}_1(\lambda))^{-1}\|.$$

Using (7.11) and (7.17) we find that (7.4) is true. ■

Proof of Proposition 7.1. The proof of (7.2) follows immediately from the Gearhart-Prüss Theorem and (7.4).

To prove (7.3) we note that since the resolvent of \mathcal{P} is compact, the spectrum is discrete. Let $\lambda_m \in \mathbb{C}$ denote an eigenvalue with real part which equals μ_m . There is clearly at least one eigenfunction associated with λ_m which we denote by u_m . We prove (7.3) by observing that

$$G(t)u_m = u_m e^{-\lambda_m t}.$$

■

APPENDIX A. SOME RESULTS RELATED TO PSEUDOSPECTRA

We recall that, for any $\epsilon > 0$, the ϵ -pseudospectra of a linear operator \mathcal{A} is defined by

$$\Sigma_\epsilon(\mathcal{A}) = \left\{ \lambda \in \mathbb{C} : \|(\mathcal{A} - \lambda)^{-1}\| > \frac{1}{\epsilon} \right\}, \quad (A.1)$$

with the convention that

$$\|(\mathcal{A} - \lambda)^{-1}\| = +\infty \quad \text{if } \lambda \in \sigma(\mathcal{A}).$$

We have

$$\bigcap_{\epsilon > 0} \Sigma_\epsilon(\mathcal{A}) = \sigma(\mathcal{A}). \quad (A.2)$$

We define, for any accretive closed operator \mathcal{A} , for $\epsilon > 0$, the ϵ -pseudospectral abscissa

$$\hat{\alpha}_\epsilon(\mathcal{A}) = \inf_{z \in \Sigma_\epsilon(\mathcal{A})} \text{Re } z. \quad (A.3)$$

We also define the growth bound of \mathcal{A} by

$$\widehat{\omega}_0(\mathcal{A}) = \lim_{t \rightarrow +\infty} \frac{1}{t} \log \|\exp(-t\mathcal{A})\|. \quad (\text{A.4})$$

We mention also that

$$\widehat{\alpha}_\epsilon(\mathcal{A}) \leq \inf_{z \in \sigma(\mathcal{A})} \operatorname{Re} z. \quad (\text{A.5})$$

Theorem A.1 (Gearhart-Prüss). *Let \mathcal{A} be a densely defined closed operator in a Hilbert space X such that $-\mathcal{A}$ generates a contraction semi-group and let $\widehat{\alpha}_\epsilon(\mathcal{A})$ and $\widehat{\omega}_0(\mathcal{A})$ denote the ϵ -pseudospectral abscissa and the growth bound of \mathcal{A} respectively. Then*

$$\lim_{\epsilon \rightarrow 0^+} \widehat{\alpha}_\epsilon(\mathcal{A}) = -\widehat{\omega}_0(\mathcal{A}). \quad (\text{A.6})$$

We refer to [11] for a proof.

REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN, *Handbook of Mathematical Functions*, Dover, 1972.
- [2] J. AGUILAR AND J. M. COMBES, *A class of analytic perturbations for one-body Schrödinger Hamiltonians*, Comm. Math. Phys., **22** (1971), 269-279.
- [3] Y. ALMOG, *The stability of the normal state of superconductors in the presence of electric currents*, SIAM J. Math. Anal., **40** (2008), 824-850.
- [4] Y. ALMOG, B. HELFFER, AND X. B. PAN, *Superconductivity near the normal state under the action of electric currents and induced magnetic fields in \mathbb{R}^2* , Commun. Math. Phys., **300** (1) (2010), 147-184.
- [5] P. BAUMAN, H. JADALLAH, AND D. PHILLIPS, *Classical solutions to the time-dependent Ginzburg-Landau equations for a bounded superconducting body in a vacuum*, J. Math. Phys., **46** (2005), art. no. 095104.
- [6] S. J. CHAPMAN, S. D. HOWISON, AND J. R. OCKENDON, *Macroscopic models for superconductivity*, SIAM Review, **34** (1992), 529-560.
- [7] J. M. COMBES AND L. THOMAS, *Asymptotic behaviour of eigenfunctions for multiparticle Schrödinger operators*, Comm. Math. Phys., **34** (1973), 251-270.
- [8] E.B. DAVIES, *Wild spectral behaviour of anharmonic oscillators*, Bull. London Math. Soc., **32** (2000), 432-438.
- [9] A. DOLGERT, T. BLUM, A. DORSEY, AND M. FOWLER, *Nucleation and growth of the superconducting phase in the presence of a current*, Phys. Rev. B, **57** (1998), 5432-5443.
- [10] Q. DU, M. D. GUNZBURGER, AND J. S. PETERSON, *Analysis and approximation of the Ginzburg-Landau model of superconductivity*, SIAM Review, **34** (1992), 54-81.
- [11] K.-J. ENGEL AND R. NAGEL, *One-parameter semigroups for linear evolution equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000.
- [12] S. FOURNAIS AND B. HELFFER, *Spectral Methods in Surface Superconductivity*, Birkhäuser, 2009.
- [13] B. HELFFER, *On spectral problems related to a time dependent model in superconductivity with electric current*, in: Proceedings of Evian conference, June 2009.
- [14] B. I. IVLEV AND N. B. KOPNIN, *Electric currents and resistive states in thin superconductors*, Advances in Physics, **33** (1984), 47-114.
- [15] T. KATO, *Perturbation Theory for Linear Operators*, Springer, third edition, 1980.
- [16] M. REED AND B. SIMON, *Methods of Modern Mathematical Physics I, Functional Analysis*, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, second edition, 1980.
- [17] J. RUBINSTEIN AND P. STERNBERG, *Formation and stability of phase slip centers in nonuniform wires with currents*, Physica C-Superconductivity and Its Applications, **468** (2008), 260-263.
- [18] J. RUBINSTEIN, P. STERNBERG, AND Q. MA, *Bifurcation diagram and pattern formation of phase slip centers in superconducting wires driven with electric currents*, Phys. Rev. Lett., **99** (2007), art. no. 167003.
- [19] J. RUBINSTEIN, P. STERNBERG, AND K. ZUMBRUN, *The resistive state in a superconducting wire: bifurcation from the normal state*, Arch. Rational Mech. Anal., **195** (2010), 117-158.
- [20] B. SIMON, *Schrödinger semigroups*, Bull. Amer. Math. Soc. (N.S.), **7** (1982), 447-526.
- [21] A. G. SIVAKOV, A. M. GLUKHOV, A. N. OMELYANCHOUK, Y. KOVAL, P. MÜLLER, AND A. V. USTINOV, *Josephson behavior of phase-slip lines in wide superconducting strips*, Phys. Rev. Lett., **91** (2003), art. no. 267001.
- [22] M. TINKHAM, *Introduction to Superconductivity*, McGraw-Hill, 1996.

- [23] D. Y. VODOLAZOV, F. M. PEETERS, L. PIRAUX, S. MATEFI-TEMPFLI, AND S. MICHOTTE, *Current-voltage characteristics of quasi-one-dimensional superconductors: An s-shaped curve in the constant voltage regime*, Phys. Rev. Lett., **91** (2003), art. no. 157001.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA.

E-mail address: `almog@math.lsu.edu`

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ PARIS-SUD 11 ET CNRS, BÂT 425, 91 405 ORSAY CEDEX, FRANCE.

E-mail address: `Bernard.Helffer@math.u-psud.fr`

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, P.R. CHINA.

E-mail address: `xbpan@math.ecnu.edu.cn`